



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


(51) International Patent Classification <sup>6</sup> : <b>C07C</b>		A2	(11) International Publication Number: <b>WO 97/07081</b>
			(43) International Publication Date: 27 February 1997 (27.02.97)
(21) International Application Number: PCT/IB96/00987		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 9 August 1996 (09.08.96)			
(30) Priority Data: 60/002,164 11 August 1995 (11.08.95) US			
(71) Applicant (for all designated States except US): YALE UNIVERSITY [US/US]; Yale Station, New Haven, CT 06520 (US).			
(72) Inventors; and		Published	
(75) Inventors/Applicants (for US only): WOOD, John, L. [US/US]; 72 Pickwick Road, Hamden, CT 06517 (US). STOLTZ, Brian, M. [US/US]; Apartment No. 1, 7 Cottage Street, New Haven, CT 06510 (US). DIETRICH, Hans-Jürgen [DE/DE]; Breslauer Strasse 31, D-47829 Krefeld-Gartenstadt (DE). PFLUM, Derek, A. [US/US]; 69 Pearl Street, New Haven, CT 06510 (US).		Without international search report and to be republished upon receipt of that report.	
(74) Agent: KRINSKY, Mary, M.; St. Onge Steward Johnston & Reens, 986 Bedford Street, Stamford, CT 06905 (US).			

(54) Title: GLYCOSYLATED INDOLOCARBAZOLE SYNTHESIS


## (57) Abstract

Tertiary alcohols containing the structural features illustrated in 3 or 4 (Scheme I) are prepared by reacting at least one diazo carbonyl compound, (e.g., 1 in Scheme I) and at least one allylic alcohol (e.g., 2 in Scheme I) in a coupling reaction run under conditions that produce carbene or carbenoid intermediates from the diazo-containing substrate such as transition metal catalysis or either thermal or photochemical decomposition. In some preferred embodiments,  $\text{Rh}_2(\text{OAc})_4$  is employed to catalyze the coupling reaction. In Scheme I R represents a substituent comprised of any number and combination of the elements H, C, N, S, Si, O, Cl, Br, I, and F. Indolocarbazoles (e.g., 7 in Scheme II) are prepared by coupling of diazo carbonyl compounds (e.g., 5 Scheme II) and biindoles (e.g., 6 in Scheme II). Indolocarbazoles are furanosylated (e.g., 7 in Scheme II) with acetals (e.g., 8 in Scheme II) or their open chain congeners (e.g., 9 in Scheme II) under conditions known to promote acetal exchange or formation, such as protic or Lewis acids. Furanosylated indolocarbazoles (e.g., 10 in Scheme II) are also prepared via ring contraction of pyranosylated indolocarbazoles (e.g., 11 in Scheme II) under conditions known to effect oxidation and benzylic acid type rearrangements, and pyranosylated indolocarbazoles (e.g., 11 in Scheme II) are prepared via ring expansion of the furanosylated congeners (e.g., 10 in Scheme II).

## Scheme I



## Scheme II



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## GLYCOSYLATED INDOLOCARBAZOLE SYNTHESIS

### **Related Application Data**

This is a continuation-in-part of U.S. provisional application serial number 60/002,164 filed August 11, 1995, which is incorporated herein by reference.

### **5 Technical Field**

This invention relates to the synthesis of tertiary alcohols by coupling of diazo carbonyl compounds with allylic alcohols under conditions that produce carbene or carbenoid intermediates. Both furanosylated and pyranosylated indolocarbazoles are prepared, including naturally occurring  
10 compounds as well as a range of structurally diverse analogues.

### **Background of the Invention**

Originally discovered in the course of screening for microbial alkaloids, staurosporine and structurally related compounds have been the object of considerable investigation for various biomedical purposes for the  
15 past twenty years (for a review, see Omura, *et al.*). It has been recently reported that staurosporine and its derivatives, for example, inhibit smooth muscle contraction, platelet aggregation, neurotrophic activity, and, most importantly, protein kinases *in vitro* and *in vivo* (*ibid.*).

Disruption of cellular signal transduction via kinase malfunction has  
20 been related to the onset of several disease states, including rheumatoid arthritis, systemic lupus erythematosus, diabetes mellitus and Alzheimer's disease. For example, the clinical severity of Alzheimer's disease correlates well with the formation of amyloid plaques and neurofibrillary tangles; both manifest paired helical filaments (PHF) that possess an  
25 overphosphorylated microtubule associated protein (M.A.P., also known as  $\tau$ -protein). It has been suggested that overphosphorylation may lead to

-2-

conformational changes that inhibit  $\tau$  binding to microtubules. Recently, a bovine  $\tau$ -kinase denominated PK40 (molecular weight 40,000) has been isolated and shown to induce a gel mobility shift of PHF- $\tau$ . PK40 is not closely associated with the cytoskeleton and appears to be a member of the extracellular regulated kinases. Specific inhibition of enzymes like PK40 by small, orally bioavailable compounds, promise to be a highly successful means of treating Alzheimer's disease.

Unfortunately, the structural homology shared by the many kinase isozymes has impeded the development of selective and therapeutically useful inhibitors. It would be desirable to have others.


### Summary of the Invention

It is a specific object of the invention to provide a synthesis for (+)- and (-)- K252a, analogues of K252a, staurosporine and its congeners, and the like.

It is another and more general object of the invention to provide for the synthesis of furanosylated and pyranosylated indolocarbzoles, particularly the interconversion of furanosylated indolocarbzoles to the corresponding pyranosylated derivatives.


It is a further object of the invention to provide an efficient approach to the synthesis of enantioenriched tertiary alcohols.

These and other objects are accomplished by the present invention, which provides a process for the preparation of tertiary alcohols containing the structural features illustrated in 3 or 4 below (Scheme I). The process utilizes at least one carbonyl compound, *e.g.*, 1 in Scheme I) and at least one allylic alcohol (*e.g.*, 2 in Scheme I) in a coupling reaction that is run under conditions that produce carbene or carbenoid intermediates from the diazo-containing substrate. These conditions include transition metal catalysis or either thermal or photochemical decomposition. In some preferred embodiments illustrated hereafter,  $\text{Rh}_2(\text{OAc})_4$  is employed to catalyze the coupling reaction.

**Schem I**

Wherein R represents a substituent comprised of any number and combination of the elements H, C, N, S, Si, O, Cl, Br, I, and F

The invention more specifically provides a process for the construction of indolocarbazoles (*e.g.*, 7 below) from the coupling of diazo carbonyl compounds (*e.g.*, 5) and biindoles (*e.g.*, 6). The invention also provides a process for the stereoselective preparation of glycosylated indolocarbazoles like but not limited to 10 and 11 via furanosylation of indolocarbazoles (*e.g.*, 7) with acetals (*e.g.*, 8) or their open chain congeners (*e.g.*, 9) under conditions known to promote acetal exchange or formation, such as protic or Lewis acids.

**Scheme II**

15

As illustrated in Scheme II above, processes of the invention also provide furanosylated indolocarbazoles (*e.g.*, 10) via ring contraction of pyranosylated indolocarbazoles (*e.g.*, 12) under conditions known to effect oxidation and benzylic acid type rearrangements. The invention correspondingly provides processes for the construction of pyranosylated indolocarbazoles (*e.g.*, 12) via ring expansion of the furanosylated congeners (*e.g.*, 10).

20


## Detailed Description of the Invention

This invention is based upon several new processes that when combined in-total or in-part can lead to the enantioselective syntheses of various indolocarbazoles.

- 5 As summarized above, in a practice of the invention at least one diazo carbonyl compound and at least one allylic alcohol of structures 1 and 2, respectively (Scheme III), are combined to produce tertiary alcohols of structures 3 and/or 4 in Scheme III. A preferred embodiment employs but is not limited to the use of transition metal catalysts in the form of ligated
- 10 Rh(II) complexes, for example  $\text{Rh}_2(\text{OAc})_4$ , to produce 3 and a Lewis acid like but not limited to  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  to convert 3 to 4. In alternative embodiments the decomposition of the diazo substrate to the corresponding carbene or carbenoid involves catalysis by complexes of: Cu(II), Mn(II), Fe(II), Co(II), Ni(0), Ni(II), Zn(II), Mo(II), Ru(II), Ru(III), Bronsted and
- 15 Lewis acids, thermolysis, and/or photolysis. The derived tertiary alcohols of structure 4 are further manipulated by standard chemical procedures to produce acetals of structure 8 and the corresponding open chain congeners of structure 9. The later are utilized in the furanosylation process and total syntheses described below.

20


**Scheme III**



(R represents a substituent comprised of any number and combination of the elements H, C, N, S, Si, O, Cl, Br, I, F)  
(X represents a substituent comprised of any number and combination of the elements N, S, Si, O, Cl, Br, I, F)

The invention also provides a process for coupling at least one diazo carbonyl compound and at least one biindole, of structures 5 and 6 respectively (Scheme IV), to produce indolocarbazoles of structure 7. A preferred embodiment employs but is not limited to the use of transition metal catalysts in the form of ligated Rh(II) complexes, for example  $\text{Rh}_2(\text{OAc})_4$ , in a solvent capable of solvating the substrates such as  $\text{CH}_2\text{Cl}_2$ , pinacolone, and/or  $\text{CH}_3\text{CN}$ . The reaction is carried out under conditions such that products are formed at a convenient rate such as for example about 20-30 minutes at reflux. In alternative embodiments initiating the process via decomposition of the diazo substrate to the corresponding carbene or carbenoid involves catalysis by complexes of: Cu(II), Mn(II), Fe(II), Co(II), Ni(0), Ni(II), Zn(II), Mo(II), Ru(II), Ru(III), Bronsted and Lewis acids, thermolysis, and/or photolysis.


#### 15 Scheme IV



(R represents a substituent comprised of any number and combination of the elements H, C, N, S, Si, O, Cl, Br, I, F)

The invention also provides a process for the stereoselective furanosylation of indolocarbazoles of a structure 10 or 11 with acetals and/or their open chain congeners, of structures 8 and 9 respectively in Scheme V, under conditions that promote acetal exchange or formation, such as but not limited to Bronstead or Lewis acids such as CSA, PTSA, or  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  (McCombie *et al.*). A preferred embodiment employs but is not limited to the use of camphor sulfonic acid (CSA) as the catalyst and the dichloroethane as the solvent in a coupling reaction that stereoselectively produces the regioisomeric furanosylated indolocarbazoles 10 and 11 in about 80% yield. The derived indolocarbazoles of structure 10 are manipulated by standard chemical procedures to produce 14 (K252a).

## Sch m V




(R represents a substituent comprised of any number and combination of the elements H, C, N, S, Si, O, Cl, Br, I, F)  
 (X represents a substituent comprised of any number and combination of the elements N, S, Si, O, Cl, Br, I, F)

As illustrated in Scheme VI, the invention also provides a process for the ring contraction of pyranosylated indolocarbazoles of structure 15 and/or 16 to furanosylated indolocarbazoles of structure 10 and/or 11 under conditions that in single- or two-step fashion can effect oxidation and benzylic acid type rearrangement (Fredenhagen *et al.*). A preferred embodiment employs but is not limited to a single-step procedure wherein CuCl is used as both the oxidant and rearrangement promoter and methanol is used as the solvent.




-7-

**Scheme VI**


(R represents a substituent comprised of any number and combination of the elements H, C, N, S, Si, O, Cl, Br, I, F)

The invention correspondingly provides a process for the ring expansion of furanosylated indolocarbazoles of structure 10 and/or 11 to the pyranosylated congeners of structure 15 and/or 16 illustrated in Scheme VII below (Ootsuka *et al.*). A preferred embodiment employs but is not limited to a multistep procedure wherein 10 and/or 11 is first reduced with  $\text{LiBH}_4$  and then the derived diol is oxidized. The resulting intermediate compound is then subjected to  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  to promote the ring expansion.

**Scheme VII**

(R represents a substituent comprised of any number and combination of the elements H, C, N, S, Si, O, Cl, Br, I, F)

It is an advantage of the invention that the combined processes provide efficient access to useful indolocarbazoles such as (+)- and (-)-K252a, (+)- and (-)-RK-286c, (+)- and (-)-MLR-52, (+)- and (-)-staurosporine, and the like depicted below.



## Examples

- 5            The following are presented to further illustrate and explain the present invention and should not be taken as limiting in any regard. Unless otherwise indicated, all parts and percentages are by weight and are based on the weight of the components at the indicated stage of processing. Rotations on indolocarbazole were obtained on methanol solutions. Compound structural assignments were in accord with infrared and high-
- 10          field  $^1\text{H}$  (500 MHz) and  $^{13}\text{C}$  (125 or 62.5 MHz) NMR spectra, as well as appropriate parent identification by high-resolution mass spectrometry.

### Example 1

#### Enantioselective Preparation of Tertiary Alcohols


- 15          This example describes a novel rhodium-catalyzed C-C bond forming reaction that allows asymmetric access to **21** (95% ee) and **22** (93% ee) in only two and three steps from methyl acetoacetate (**18**) (Scheme VIII). In this scenario  $\alpha$ -keto ester **21** was produced from the rhodium-catalyzed decomposition of **19** in the presence of S-(+)-1-buten-3-ol (**20**) (Wood *et al.*). In the event, complete consumption of **19** was observed after only 20

-9-

minutes to reflux in benzene. Proton NMR analysis of the crude reaction indicated the clean

formation of a product similar to 19; however, the characteristic methyl ketone singlet had shifted from 2.2 to 1.5 ppm. Clearly the allyloxy or allyloxonium ylide intermediate had undergone [3,3] sigmatropic rearrangement to alcohol (+)-21 (66% yield) (Pirrung *et al.*). Completion of the tandem rearrangement protocol was achieved by exposing (+)-21 to  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  which promoted a clean [1,2]-allyl migration to furnish (-)-22 in 74% yield. In subsequent studies, improved yields were obtained by conducting the tandem rearrangement in one pot. Thus, introducing an equivalent of  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  into the cooled [3,3] reaction allows isolation of (-)-22 in an overall yield of 75%.


### Scheme VIII



With an approach firmly established, a chemical correlation study was initiated to confirm both the sense and degree of asymmetric induction for the tandem rearrangement. Analysis of the purified products from both the [3,3] (*i.e.*, (+)-21) and [1,2] (*i.e.*, (-)-22) rearrangements via proton NMR in the presence of  $\text{Eu}(\text{hfc})_4$  gave the first indication that each step was proceeding with a high degree of stereoselectivity. Conversion of (+)-21 to 23 as outlined in Scheme IX, followed by comparison of the derived bis Mosher ester (24) to samples prepared from *S*-(+)- and *R*-(-)-citramalic acid, established that *S*-(+)-20 (98% ee) had furnished *R*-(+)-21 (95% ee). Stereoselectivity in the [1,2] shift was established by degradation of (-)-22 to *R*-(-)-25 followed by DIBAL reduction and proton NMR analysis of the corresponding bis Mosher ester. While the Mosher ester analysis established an ee of 92%, the observation of *R*-(-)-25 in the degradation proved the absolute stereochemistry in 22 as *S*.


-10-

## Scheme IX



Having established the degree and sense of asymmetric induction the asymmetric synthesis of the requisite acetals **26** and **27** was begun. Thus, reductive ozonolysis of **22** followed by acetal formation provided a ternary mixture. Spectral identification of the isolated products indicated the presence of methyl ketone **26** and furanoses (+)-**27a** and (+)-**27b** (Scheme X).

## Scheme X




## Example 2

Preparation and Furanosylation of Indolocarbazoles  
The Synthesis of K252a

This example describes the coupling of diazolactams **28** and 2,2'-biindole **29** to produce an intermediate that undergoes cycloaromatization to furnish the indolocarbazoles **30**. Application of this strategy allows efficient access to both the parent aglycone (**30a**) and the selectively protected derivatives (**30b-c**). Of the latter, **30c** is employed in the total synthesis of K252a. Overall, preparation of the enantioenriched furanoses **27** (described in Example 1 above) and aglycon unit **30c** and their

conversion to **14** require only eleven synthetic operations with a longest linear sequence of seven steps.

### Scheme XI




- 5           The feasibility of the carbenoid approach to **30** was initially assessed by reaction of **28a** (1.0 equiv) with indole (3.0 equiv) in the presence of catalytic  $\text{Rh}_2(\text{OAc})_4$  (0.01 equiv, Scheme XII). After only 12 h, TLC analysis indicated complete consumption of **28a** and standard work-up and isolation procedures furnished **31** in 65% yield. Similar conditions proved
- 10 ineffective for the coupling of **28a** with **29**, and it was only after considerable experimentation that a procedure was developed which provided satisfactory yields of **30a**. The use of degassed pinacolone proved critical as this solvent was found to be both compatible with the carbenoid chemistry and capable of solvating the diindole substrate. Under these conditions the
- 15 coupling of **28a** and **29** proceeded directly to **30a** (K252c) in 25% yield. Presumed intermediates **32** and **33** were not apparent by TLC or NMR analysis of the crude reaction mixture. In an attempt to complete the synthesis, the cycloglycosidation of **27** with **30a** revealed a tendency of the latter to alkylate at the amide nitrogen; thus, selectively protected
- 20 aglycones **30b-e** were employed. Preparation of the corresponding diazotactams **28b-e**, followed by reaction with **29** in the presence of  $\text{Rh}_2(\text{OAc})_4$  (0.1 equiv) established that several protecting groups can withstand the carbenoid conditions and that the best yields (50-62%) are obtained within the benzyl class (*e.g.*, **28c,d,e**→**30c,d,e** Scheme XII). To

-12-


provid th most flexibility in the eventual deprotection 3,4-dimethoxybenzyl protected aglycone **28** was employed.

## Sch m XII



- 5 Having gained efficient access to **30c**, attention was turned to the preparation of the furanose components (**27**). To this end, a novel tandem rearrangement protocol was developed that combines methyl 2-diazo-3-oxobutyrates (**19**) and *S*-(+)-1-buten-3-ol (**20**) to furnish (-)-**22** in a single-pot (92% ee, 75% yield). Reductive ozonolysis of (-)-**22** followed by acid promoted cyclization in methanol produced (+)-**27a** and (+)-**27b** in good yield.
- 10

## Scheme XIII




- 15 With both (+)-**27** and **30c** in hand the cycloglycosidative coupling was investigated. Of several conditions reported by McCombie, *et al.*, for related transformations, camphorsulfonic acid in 1,2-dichloroethane was found to be th catalyst and solvent of choice. In th vent, **30c** and (+)-**27a** and

-13-

**27b** combined rapidly to form two regioisomeric pairs of open chain monoamino acetal diastereomers (**34** and **35**). Prolonged heating of the quaternary mixture induced cycloglycosidation to only two of the four possible diastereomers.

5 **Scheme XIV**




Preliminary assignment of structure was based on <sup>1</sup>H NMR analysis which indicated that the reaction had produced the regioisomeric products **(-)-36** (55% yield) and **(-)-37** (25% yield). The observed formation of **(-)-14** upon deprotection of **(-)-36** under standard conditions (TFA/CH<sub>2</sub>Cl<sub>2</sub>/thioanisole) established the cycloglycosidation as both regio- and stereoselective for the natural configuration. Comparison of synthetic

(-)-14 to material derived from natural sources established its identity as the unnatural enantiomer of K252a.

Total synthesis of the natural enantiomer (*i.e.*, (+)-14) was effected in an analogous fashion using 28c and (-)-27a and (-)-27b as coupling partners (Scheme XV). The latter compound was prepared via the tandem [3,3]/[1,2] rearrangement protocol (described in Example 1 above) using *R*-(-)-1-nonen-3-ol (38) as the source of asymmetry.

#### Scheme XV



In summary, application of a novel carbenoid mediated synthesis of K252c coupled with a highly selective tandem (3,3)/(1,2) rearrangement protocol provides efficient access to both (+)- and (-)-K252a.

#### Example 3

Furanosylation of Alternative Indolocarbazoles and the Interconversion of Furanosylated and Pyranosylated Indolocarbazoles

This example reports results wherein an indolocarbazole simpler than that described Example 2 is subjected to furanosylation. The derived product 40 is further manipulated into a ring-expansion substrate 41 or 42 that undergoes conversion to the corresponding pyranosylated indolocarbazole 43 or 44, respectively (Stoltz *et al.* 1995). It is further demonstrated in this example that the  $\alpha$ -hydroxy ketone congener 43 undergoes facile oxidative ring contraction to the furanosylated indolocarbazole upon exposure to CuCl in methanol (Stoltz *et al.* 1996).


For the furanosylation, indolo [2,3-*a*]carbazole (39) was coupled with 27a and 27b in a manner similar to that employed in the synthesis of



K252a described in Example 2. This coupling again proved highly stereoselective and produced **40** as the only isolable product in 85% yield.

### Scheme XVI


5



Turning to the ring expansion, it was soon discovered that transformation of **40** into aldehyde **41** followed by treatment with BF<sub>3</sub>•Et<sub>2</sub>O results in a regio- and stereoselective rearrangement to the pyranosylated indolocarbazole **43**. At this stage all that remained for the preparation of **44** was what appeared to be a trivial alkylation of the C(3') hydroxyl. Ketone **43** surprisingly proved quite resistant to methylation under numerous alkylation conditions. In addition, attempts to incorporate directly the methyl substituent by promoting the rearrangement with a source of Me<sup>+</sup> (e.g., Meerwein's reagent, TMSOTf/TMSOMe, and MeOTf) also failed. Eventually, these difficulties led to the development of an alternative strategy that targeted dimethyl acetal **42** as the substrate for a ring expansion (Scheme XVII). Although **42** was readily produced under a variety of conditions, its instability to chromatographic purification required the employment of montmorillonite clay K-10 to promote acetal formation. Removal of the clay via filtration, solvent exchange with Et<sub>2</sub>O, and subsequent treatment with BF<sub>3</sub>•Et<sub>2</sub>O resulted in the slow (72 h, 25 °C) conversion of **42** to **44** (50% yield).


-16-

## Scheme XVII



Having rapidly assembled  $\alpha$ -methoxy ketone 44, its conversion to the desamido pyranosylated indolocarbazoles was investigated. To this end, the analogs of RK-286c (46) and TAN-1030a (47) were readily prepared from 44 under standard conditions using  $\text{NaBH}_4$  and  $\text{H}_2\text{NOH} \cdot \text{HCl}$ , respectively (Scheme XVIII).

## Scheme XVIII




While the above model investigations established the feasibility of a regio and stereoselective ring expansion, subsequent attempts to alkylate the derived  $\alpha$ -hydroxy ketone 43 proved problematic. Of note is the propensity of 43 to undergo loss of the indolocarbazole nucleus as evidenced

-17-

by isolation of **39** as the major product in many of the alkylation attempts. In an effort to avoid this deleterious event attention was turned to methylation procedures that appeared to proceed under essentially neutral conditions. While these efforts failed to produce any of the desired  $\alpha$ -methoxy ketone **44**, the conditions comprising CuCl and DCC in MeOH were observed to cleanly convert **43** to **40**, the functionalized K252a sugar moiety. Apparently these conditions induced either ring contractive  $\alpha$ -ketol rearrangement and oxidation (*i.e.*, **43**→**41**→**40**) or oxidation and ring contractive "benzilic" acid rearrangement (*i.e.*, **43**→**48**→**40**). While not wishing to be bound to any theory, since  $\alpha$ -hydroxy aldehyde **41** failed to undergo conversion to **40** under identical conditions, the latter of these two mechanistic possibilities appears most likely. In addition, subsequent investigations have revealed CuCl in MeOH without added DCC to be the optimal conditions for converting **43** to **40** (95% yield).

#### Scheme XIX




#### Example 4

##### The Synthesis of Staurosporine, RK-286c, MLR-52, and K252a

This example demonstrates that the tertiary alcohol and indolocarbazole syntheses, the indolocarbazole furanosylation, and the ring-expansion protocol described in the above Examples can be used to prepare pyranosylated indolocarbazoles that are suited for conversion to staurosporine (**49**) RK-286c (**50**), MLR-52 (**51**), and K252a (**14**) (Link *et al.*).

The synthesis of 49-51 began by converting the K252a precursor (36, described above in Example 2) to the corresponding aldehyde via  $\text{LiBH}_4$  reduction and then Moffatt oxidation (63% yield overall, Scheme XX). Guided by the  $\alpha$ -ketol rearrangement results described above, 50 was exposed to  $\text{BF}_3 \cdot \text{Et}_2\text{O}$  and the reaction allowed to stir at room temperature for 3 h. Given that the proposed ring-expansion of 50 to 51 could proceed to a mixture of regio- and stereoisomeric products, treatment of (+)-50 with  $\text{BF}_3 \cdot \text{OEt}_2$  in  $\text{Et}_2\text{O}$  (2.2 equiv, 25- 30 °C, 24h) surprisingly produces a single product, (+)-51, in 85% yield. The regio- and stereochemical outcome of this reaction, which were confirmed by spectral comparison to a closely related model and the conversion of (+)-51 to (+)-50 (*vide infra*).

#### Scheme XX




As expected from the data presented in Example 3, attempts to methylate 51 were unproductive and again led to the observation that exposure of (+)-51 to  $\text{CuCl}$  in  $\text{MeOH}$  results in a highly stereoselective oxidation/ring-contraction sequence that produces (+)-36 in 95% yield.

Turning from the potentially biomimetic synthesis of (+)-K252a to the synthesis of 49-51, it was discovered that (+)-51 undergoes selective conversion to (+)-54 upon sequential treatment with  $\text{NaBH}_4$  and  $\text{NaH}/\text{MeI}$ . Having installed all of the functional groups common to (+)-50-51, the


approach diverged into the synthesis of (+)-RK286c and (+)-MLR-52. The former was completed via deprotection of (+)-54 (TFA/anisole) while the latter required a three-step sequence that was initiated by exposing (+)-54 to Martin's sulfurane. Oxidation of the derived olefin with OsO<sub>4</sub> followed by deprotection of the resultant diol (+)-55 produced (+)-51 (Scheme XXI).

### Scheme XXI



The inability to prepare  $\alpha$ -methoxy ketone 52 guided an approach to staurosporine along a route wherein the 4' nitrogen is introduced via conversion of (+)-51 to the corresponding oxime (-)-56 (H<sub>2</sub>NOH·HCl, NaOAc, Scheme XXII). Crucial for the success of this approach is the fact that (-)-56, unlike ketone (+)-51, readily undergoes alkylation to the C(3') methyl ether (MeI, KOH, *n*-Bu<sub>4</sub>NBr). Stereoselective reduction of the derived methoxy oxime (-)-57 (H<sub>2</sub>, PtO<sub>2</sub>) to the corresponding primary amine ((+)-58) followed by monomethylation (HCO<sub>2</sub>COCH<sub>3</sub>, BH<sub>3</sub>·DMS) and deprotection (TFA) produced (+)-staurosporine (49).

## Schem XXII



In summary, efforts to devise an efficient synthesis of the pyranosylated indolocarbazoles via a common intermediate [*i.e.*, (+)-36] were successful in delivering (+)-49 (19 steps), (+)-50 (17 steps), and (+)-51 (19 steps). In addition, these investigations have revealed both ring-expansion and -contraction reactivity that may play a central role in the biogenesis of both the furanosylated and pyranosylated members of this important class of natural products.

10

## Example 5

## Experimental Procedures for Selected Compounds in Examples 1-4

Preparation of 21. (R)-3-Hydroxy-3-methyl-2-oxo-5-heptenoic acid ethylester 2.1323 g (15.00 mmol) 2-diazomethylacetoacetate are dissolved in 75 ml benzene (1 neck 100 ml flask). Under stirring 1.0816 g (15 mmol; 1.3 ml) (S)-(+)-butenol are added to the solution. After addition of 66.3 mg (0.15 mmol; 0.01 eq.) Rh<sub>2</sub>(OAc)<sub>4</sub> the flask is immersed into a preheated (100-110 °C) oil bath. The mixture is heated under reflux for 30 minutes. After 1 minute vigorous nitrogen evolution starts and lasts for about 2 minutes. After cooling the mixture the solvent is evaporated and the residue is

- flashed on silica (4.5 x 15 cm) using hexane/ethylacetate (20%). 1.66 g (59%) of the product are obtained. This relatively low yield (compared to the 74% obtained in the racemic series) must be due to same impurity (water?) in the butenol. b.p. 65-67°/0.35 mm Hg; IR (thin film/NaCl) 3521.0 (m), 3028.5 (w), 2981.5 (m), 2957.1 (m), 2937.9 (m), 2919.9 (m), 2857.4 (w), 1742.6 (s), 1726.1 (s), 1452.3 (m), 1437.5 (m), 1376.0 (w), 1361.2 (w), 1289.1 (m), 1250.1 (m), 1192.6 (w), 1145.8 (w), 1116.3 (w), 1081.5 (w), 1060.1 (w), 1032.1 (s), 971.9 (m), 920.3 (w), 861.6 (w), 844.7 (w), 814.4 (w), 722.7 (w), 663.1 (w)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  5.57 (m, 1H), 5.35 (m, 1H), 3.88 (s, 3H), 3.28 (br.s, 1H), 2.68 (dd,  $J=7.0, 14.0$  Hz, 1H), 2.42 (dd,  $J=7.7, 14.0$  Hz, 1H), 1.66 (d,  $J=6.42$  Hz, 3H), 1.47 (s, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  198.58, 162.78, 130.97, 123.58, 78.30, 52.55, 42.26, 24.15, 17.82; HRMS (CI, isobutane)  $m/z$  calc'd for  $\text{C}_9\text{H}_{15}\text{O}_4$  (M+H): 187.0970, found 187.0966;  $[\alpha]_D^{22} + 14.65^\circ$  ( $c=1.08$ ,  $\text{CHCl}_3$ ).
- 15 Preparation of 22. (S)-2-Hydroxy-2-allylmethylmethylacetoacetate 3.35 g (18 mmol) (R)-3-hydroxy-3-methyl-2-oxo-5-heptenoicacidethyl-ester are dissolved in 180 ml dichloromethane and treated with 2.554 g (18 mmol; 2.21 ml)  $\text{BF}_3 \cdot \text{OEt}_2$ . The reaction mixture is stirred 2 hours at 25 °C (TLC control). The solvent is evaporated and the residue flashed on silica (4.5 x 18 cm) using hexane/ethylacetate (20%). 2.368 g (71%) of the product are isolated. IR (thin film/NaCl) 3476.1 (m), 3031.2 (w), 3009.6 (w), 2956.2 (m), 2921.4 (w), 2857.5 (w), 1746.9 (s), 1721.9 (s), 1437.4 (m), 1357.9 (m), 1271.0 (m), 1224.2 (m), 1195.9 (m), 1183.2 (m), 1141.0 (m), 1108.5 (m), 1076.9 (w), 1052.8 (w), 994.6 (w), 972.4 (m), 861.8 (w), 816.7 (w), 798.3 (w)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  5.60 (m, 1H), 5.32 (m, 1H), 4.17 (s, 1H), 3.80 (s, 3H), 2.77 (dd,  $J=6.6, 14.3$  Hz, 1H), 2.63 (dd,  $J=7.6, 14.3$  Hz, 1H), 2.28 (s, 3H), 1.65 (d,  $J=6.5$  Hz, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  204.28, 170.85, 130.59, 122.91, 83.89, 53.16, 38.54, 24.76, 17.94; HRMS (CI, isobutane)  $m/z$  calc'd for  $\text{C}_9\text{H}_{15}\text{O}_4$  (M+H): 187.0970, found 187.0969;  $[\alpha]_D^{22} - 32.13^\circ$  ( $c=1.08$ ,  $\text{CHCl}_3$ ).
- 20 (S)-2-Hydroxy-2-allylmethylmethylacetoacetate (1 pot procedure) 426.5 mg (3.00 mmol) 2-diazomethylacetoacetate are dissolved in 15 ml benzene (1 neck 25 ml flask). Under stirring 237.9 mg (3.3 mmol; 0.286 ml; 1.1 eq.) (S)-(+)-butenol are added to the solution. After addition of 13.3 mg (0.03 mmol; 0.01 eq.)  $\text{Rh}_2(\text{OAc})_4$  the flask is immersed into a preheated (100-110 °C) oil bath. The mixture is heated under reflux for 30 minutes. After 1 minute vigorous

-22-

nitrogen evolution starts and lasts for about 2 minutes. After cooling the reaction mixture 554.5 mg (3.75 mmol; 0.46 ml; 1.25 eq.)  $\text{BF}_3 \cdot \text{OEt}_2$  are added. The mixture is stirred for 2-3 hours (TLC control) at 25 °C. The reaction mixture is passed through silica (2.5 x 8 cm) using pentane/ether (20%) as solvent; 416.9 mg (75%) of the product are isolated.

Preparation of 27. (S)-(+)-Hydroxy furanose and (S)-(-)-methylketonedimethylacetal 1.305 g (7.00 mmol) (S)-2-hydroxy-2-allylmethylmethylacetoacetate and a trace of sudan red dye are dissolved in 45 ml methanol. After cooling to -78 °C the mixture is treated with  $\text{O}_3$  until the dye is completely discolored (about 3 minutes). The mixture is purged with argon for 10 Minutes at -78 °C and 20 ml dimethylsulfide are added at that temperature. The dry ice cold bath is replaced with an ice bath which is allowed to thaw (0-20 °C) over a period of 3 hours. The solvent is removed and the crude product dissolved in 45 ml benzene. After addition of 20.0 mg (0.105 mmol; 0.015 eq.) p-toluenesulfonic acid and 12 ml methanol the mixture is stirred at 25 °C for 17 hours (until the reaction is completed as judged by TLC). The solvent is evaporated and the product is flashed on silica (4.5 x 20 cm) using hexane/ethylacetate (20%) as solvent system. 1.69 g (80%) of a mixture (1:1:1) of two furanose diastereomers and the methylketonedimethylacetal is obtained. They can be separated using HPLC. In a first run (stationary phase:  $\text{SiO}_2$ ; mobile phase: hexane/dichloromethane/ethylacetate (2:2:1) a mixture of the furanose diastereomer I and the methylketone is eluted first followed by the second furanose diastereomer which can be isolated in its pure form. The two component mixture is separated using a different system (stationary phase:  $\text{SiO}_2$ ; mobile phase: hexane/i-propanol (10%)). The furanose diastereomer I is eluted as first fraction closely followed by the methylketonedimethyl acetal. Hydroxyfuranose mp 81-82 °C; IR (thin film/ $\text{NaCl}$ ) 3495.1 (m), 2995.3 (m), 2953.2 (s), 2917.2 (s), 2848.3 (m), 1747.11 (s), 1463.77 (m), 1439.35 (m), 1379.1 (m), 1355.2 (w), 1263.5 (s), 1200.0 (s), 1182.1 (m), 1156.1 (m), 1121.86 (s), 1086.1 (s), 1043.7 (m), 1019.3 (m), 973.677 (m), 949.3 (m), 929.7 (m), 892.27 (m), 864.6 (w), 833.7 (m), 802.7 (m), 750.6 (m), 685.5 (m)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  5.07 (dd,  $J=0.6, 5.8$  Hz, 1H), 3.79 (s, 3H), 3.42 (s, 3H), 3.38 (br. s, 1H), 3.25 (s, 3H), 3.03 (dd,  $J=5.8, 14.2$  Hz, 1H), 2.06 (d,  $J=14.2$  Hz, 1H), 1.54 (s, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  170.50, 110.55, 103.86, 83.17, 55.56, 52.63,



- 49.28, 40.56, 15.82;  $[\alpha]_D^{20} + 112.13^\circ$  ( $c=1.06$ ,  $\text{CHCl}_3$ ). Methylketone IR (thin film/ $\text{NaCl}$ ) 3450.0 (m), 2988.3 (m), 2953.5 (s), 2915.0 (s), 2849.2 (s), 1746 (s), 1722.3 (s), 1457.5 (m), 1436.4 (m), 1386.7 (m), 1275.0 (m), 1245.2 (m), 1198.0 (m), 1178.1 (m), 1142.1 (s), 1121.0 (s), 1063.3 (s), 1014.2 (w), 998.1 (w), 974.4 (w), 907.4 (w), 830.6 (w), 755.1 (w)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  4.50 (s, 1H), 4.50 (dd,  $J=4.8, 6.7$  Hz, 1H), 3.78 (s, 3H), 3.34 (s, 3H), 3.29 (s, 3H), 2.43 (dd,  $J=4.8, 14.5$  Hz, 1H), 2.39 (dd,  $J=6.7, 14.5$  Hz, 1H), 2.28 (s, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  204.04, 170.87, 102.01, 81.80, 54.99, 53.84, 53.22, 38.40, 24.50;  $[\alpha]_D^{20} - 20.25^\circ$  ( $c=0.97$ ,  $\text{CHCl}_3$ ).
- 10 Hydroxyfuranose mp 63-64  $^\circ\text{C}$ ; IR (thin film/ $\text{NaCl}$ ) 3480.7 (m), 2995.0 (w), 2953.3 (m), 2914.2 (w), 2835.1 (w), 1726.7 (s), 1443.2 (m), 1377.9 (m), 1348.2 (w), 1278.2 (s), 1239.0 (m), 1228.0 (m), 1200.4 (m), 1181.6 (w), 1165.1 (s), 1127.6 (s), 1114.3 (s), 1092.2 (m), 1084.5 (m), 979.6 (m), 957.5 (m), 948.7 (m), 927.8 (m), 901.5 (m), 871.9 (w), 840.9 (w), 802.9 (w), 755.5 (m), 673.0 (w)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  5.21 (app. t,  $J=5.7$  Hz, 3H), 3.79 (s, 3H), 3.47 (s, 3H), 3.36 (d,  $J=1.6$  Hz, 1H), 3.27 (s, 3H), 2.84 (ddd,  $J=1.6, 5.2, 14.3$  Hz, 1H), 2.34 (dd,  $J=6.2, 14.3$  Hz, 1H), 1.43 (s, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  172.10, 1089.84, 105.38, 84.46, 56.37, 52.82, 49.00, 40.44, 14.46; HRMS (CI, isobutane)  $m/z$  calc'd for  $\text{C}_8\text{H}_{13}\text{O}_5$  (M- $\text{CH}_3\text{OH}+\text{H}$ ): 189.0763, found 189.0764;  $[\alpha]_D^{22} + 9.66^\circ$  ( $c=1.03$ ,  $\text{CHCl}_3$ ).
- 20

- Preparation of 38. (R)-2-Hydroxy-2-allylhexylmethylacetoacetate 426.5 mg (3.00 mmol) 2-diazomethylacetoacetate are dissolved in 15 ml benzene (1 neck 25 ml flask). Under stirring 469.3 mg (3.3 mmol; 1.1 eq.) (R)-(-)-nonenol are added to the solution. After addition of 13.3 mg (0.03 mmol; 0.01 eq.)  $\text{Rh}_2(\text{OAc})_4$  the flask is immersed into a preheated (100-110  $^\circ\text{C}$ ) oil bath. The mixture is heated under reflux for 30 minutes. After 1 minute vigorous nitrogen evolution starts and lasts for about 2 minutes. After cooling the reaction mixture 554.5 mg (3.75 mmol; 0.46 ml; 1.25 eq.)  $\text{BF}_3 \cdot \text{OEt}_2$  are added. The mixture is stirred for 2-3 hours (TLC control) at 25  $^\circ\text{C}$ . The reaction mixture is passed through silica (2.5 x 8 cm) using pentane/ether (20%) as solvent; 510.9 mg (66%) of the product are isolated. R)-(-)-Hydroxy furanose and (R)-(+)-methylketonedimethylacetal The same procedure that was employed for the preparation of (S)-(+)-hydroxy furanose and (S)-(-)-methylketonedimethylacetal is used. 1.798 g (7.00 mmol) of (R)-2-hydroxy-2-allylhexylmethylacetoacetate are used as
- 25
- 30
- 35

- starting material yielding 1.04 g (68%) of the 3 component mixture (1:1:1). Hydroxyfuranose mp 81-82 °C; IR (thin film/NaCl) 3496.4 (m), 2998.9 (m), 2953.3 (m), 2915.1 (m), 2836.9 (m), 1748.9 (s), 1732.9 (s), 1440.3 (m), 1379.3 (m), 1334.7 (w), 1261.7 (s), 1200.7 (s), 1182.7 (m), 1156.9 (s), 1122.5 (s), 1098.5 (s), 1086.5 (s), 1044.3 (m), 1021.1 (m), 975.9s, 94826 (m), 930.7 (m), 893.9 (m), 865.4 (m), 834.9 (m), 802.2 (m), 750.9 (m), 685.7 (m)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  5.07 (d,  $J=5.8$  Hz, 1H), 3.78 (s, 3H), 3.42 (s, 3H), 3.25 (s, 3H), 3.03 (dd,  $J=5.8, 14.1$  Hz, 1H), 2.05 (d,  $J=14.1$  Hz, 1H), 1.54 (s, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  170.41, 110.46, 103.78, 83.08, 55.47, 52.52, 49.18, 40.49, 15.73;  $[\alpha]_{\text{D}}^{20} - 122.55^\circ$  ( $c=1.10$ ,  $\text{CHCl}_3$ ). Methylketone IR (thin film/NaCl) 3452.5 (m), 2993.2 (m), 2954.6 (m), 2934.2 (m), 2917.5 (m), 2848.4 (m), 2838.2 (m), 1748.7 (s), 1723.1 (s), 1437.8 (m), 1359.7 (m), 1275.8 (m), 1245.7 (m), 1198.5 (m), 1178.2 (m), 1144.7 (s), 1124.4 (s), 1065.4 (s), 1015.7 (m), 997.3 (m), 905.2 (m), 829.6 (w), 802.0 (w), 756.0 (w)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  4.51 (br. s, 1H), 4.50 (dd,  $J=4.9, 6.6$  Hz, 1H), 3.78 (s, 3H), 3.34 (s, 3H), 3.29 (s, 3H), 2.43 (dd,  $J=4.9, 14.6$  Hz, 1H), 2.38 (dd,  $J=6.6, 14.6$  Hz, 1H), 2.28 (s, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  204.00, 170.84, 101.96, 81.76, 54.92, 53.79, 53.18, 38.37, 24.46;  $[\alpha]_{\text{D}}^{20} + 19.55^\circ$  ( $c=1.12$ ,  $\text{CHCl}_3$ ). Hydroxyfuranose mp 63-64 °C; IR (thin film/NaCl) 3486.7 (m), 2994.8 (m), 2954.8 (m), 2918.0 (m), 2836.2 (m), 1732.7 (s), 1442.6 (m), 1378.3 (m), 1346.6 (w), 1276.5 (s), 1243.0 (m), 1229.7 (m), 1199.7 (m), 1183.0 (m), 1165.4 (s), 1126.7 (s), 1115.6 (s), 1086.6 (s), 1049.2 (s), 1020.2 (s), 980.1 (m), 956.6 (m), 626.1 (m), 902.6 (m), 870.2 (w), 840.2 (w), 803.0 (w), 754.6 (m), 673.3 (m)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  5.21 (app. t,  $J=5.7$  Hz, 1H), 3.79 (s, 3H), 3.47 (s, 3H), 3.36 (br. s, 1H), 3.27 (s, 3H), 2.84 (dd,  $J=5.3, 14.3$  Hz, 1H), 2.34 (dd,  $J=6.2, 14.3$  Hz, 1H), 1.43 (s, 3H);  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ )  $\delta$  172.11, 109.86, 105.40, 84.48, 56.39, 52.83, 49.02, 40.46, 14.48;  $[\alpha]_{\text{D}}^{20} -9.00^\circ$  ( $c=1.16$ ,  $\text{CHCl}_3$ ).
- General method for the preparation of 30a-d Dry  $\text{N}_2$  is bubbled through a mixture of 2,2'-biindole (0.86 mmol), diazo compound 4 (2.2 mmol, 2.5 eq),  $\text{Rh}_2(\text{OAc})_4$  (0.086 mmol, 0.1 eq) and 8.6 mL pinacolone, in a 3-neck round bottom flask fitted with a reflux condenser for 2 h. The reaction mixture is then heated to reflux for 8 h. The mixture is allowed to cool to 25 °C, the solvent is evaporated, the residue is chromatographed (1:1 EtOAc:Hexanes) to provide (R= 3,4-DMB 0.25 g, 0.56 mmol, 65%; R=4-

PMB 0.22 g, 0.47 mmol, 55%; R=Bn 0.20 g, 0.5 mmol, 58%; R=*t*-Bu 0.13 g, 0.34 mmol, 40%).

Indolocarbazole **30a**. IR (thin film/NaCl) 3485.3 (brm), 3456.0 (brm), 3343.1 (brs), 3249.7 (brm), 2979.7 (m), 1654.4 (w), 1600.5 (s), 1578.2 (s),  
5 1465.8 (w), 1446.5 (m), 1385.0 (s), 1364.0 (m), 1335.9 (w), 1225.3 (s)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.45 (bs, 1H), 11.29 (bs, 1H), 9.24 (d,  $J=7.9$  Hz, 1H), 8.09 (d,  $J=7.8$  Hz, 1H), 7.77 (d,  $J=8.2$  Hz, 1H), 7.70 (d,  $J=8.2$  Hz, 1H), 7.47 (app.t,  $J=7.5$  Hz, 1H), 7.41 (app.t,  $J=7.5$  Hz, 1H), 7.30 (app.t,  $J=7.5$  Hz, 1H), 7.21 (app.t,  $J=7.5$  Hz, 1H), 5.13 (s, 2H), 1.65 (s, 9H);  $^{13}\text{C}$   
10 NMR (62.5 MHz, DMSO- $d_6$ )  $\delta$  169.9, 139.2, 139.0, 129.9, 127.6, 125.4, 125.3, 124.9, 122.7, 122.4, 122.0, 121.2, 119.7, 118.8, 115.1, 113.6, 111.8, 111.2, 101.9, 53.6, 48.1, 27.8; HRMS (FAB)  $m/z$  calc'd for  $\text{C}_{24}\text{H}_{22}\text{N}_3\text{O}$  (M+H): 368.1762, found 368.1764.

Indolocarbazole **30b**. IR (thin film/NaCl) 3487.5 (brs), 3352.0 (brs), 3232.0 (brs), 3022.3 (m), 1579.1 (s), 1571.2 (s), 1517.7 (s), 1462.9 (s), 1399.3 (m),  
15 1262.7 (m), 1237.6 (s), 1142.0 (w), 1016.8 (w), 741.3 (s)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.50 (bs, 1H), 11.35 (bs, 1H), 9.28 (d,  $J=7.9$  Hz, 1H), 7.97 (d,  $J=7.8$  Hz, 1H), 7.77 (d,  $J=8.1$  Hz, 1H), 7.73 (d,  $J=8.1$  Hz, 1H), 7.45 (app.t,  $J=6.9$  Hz, 1H), 7.44 (app.t,  $J=7.1$  Hz, 1H), 7.26 (app.t,  $J=7.1$  Hz, 1H), 7.25 (app.t,  $J=7.1$  Hz, 1H), 7.02 (s, 1H), 6.92 (s, 2H), 4.94 (s, 2H),  
20 4.82 (s, 2H), 3.74 (s, 3H), 3.71 (s, 3H);  $^{13}\text{C}$  NMR (62.5 MHz, DMSO- $d_6$ )  $\delta$  169.2, 148.9, 148.1, 139.1, 139.0, 130.6, 130.0, 127.7, 125.3, 124.9, 124.9, 124.8, 122.6, 122.3, 120.7, 119.9, 119.7, 118.8, 118.2, 115.4, 113.8, 112.3, 112.1, 111.7, 111.1, 55.5, 49.3, 45.4; HRMS (FAB)  $m/z$  calc'd for  $\text{C}_{29}\text{H}_{24}\text{N}_3\text{O}_3$  (M+H): 462.1817, found 462.1813.  
25

Indolocarbazole **30c**. IR (thin film/NaCl) 3429.3 (brs), 3351.3 (brs), 2912.4 (m), 1609.7 (s), 1580.3 (s), 1512.0 (s), 1465.5 (s), 1402.1 (w), 1250.61 (s),  
1238.4 (s), 1177.3 (m), 1030.8 (w), 748.9 (s)  $\text{cm}^{-1}$ ;  $^1\text{H}$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  11.53 (bs, 1H), 11.37 (bs, 1H), 9.28 (d,  $J=7.8$  Hz, 1H), 7.99 (d,  $J=7.7$  Hz, 1H), 7.78 (d,  $J=8.1$  Hz, 1H), 7.75 (d,  $J=8.1$  Hz, 1H), 7.47 (app.t,  $J=7.0$  Hz, 1H), 7.45 (app.t,  $J=7.1$  Hz, 1H), 7.36 (d,  $J=8.4$  Hz, 2H), 7.28 (app.t,  $J=7.9$  Hz, 1H), 7.26 (app.t,  $J=7.8$  Hz, 1H), 6.94 (d,  $J=8.5$  Hz, 2H),  
30 4.94 (s, 2H), 4.83 (s, 2H), 3.72 (s, 3H);  $^{13}\text{C}$  NMR (62.5 MHz, DMSO- $d_6$ )  $\delta$  169.2, 158.4, 139.1, 139.0, 130.0, 129.9, 128.9, 127.7, 125.3, 124.9, 124.8,

122.6, 122.2, 120.7, 119.7, 118.8, 118.2, 115.4, 113.9, 113.8, 111.7, 111.1, 54.9, 49.2, 45.0; HRMS (FAB)  $m/z$  calc'd for  $C_{28}H_{22}N_3O_2$  (M+H): 432.1712, found 432.1699.

(+)-N-3,4-Dimethoxybenzyl-K252a (**36**). Aglycone **30c** (0.22 mmol) and (-)-2,5-dimethoxy sugar (0.87 mmol, 4.0 eq) were refluxed in dry 1,2-dichloroethane (7.5 mL) in the presence of camphorsulfonic acid (0.022 mmol, 0.1 eq) for 48 h. The reaction mixture was allowed to cool to 25 °C, diluted with 5.0 mL  $CH_2Cl_2$ , and washed with 5.0 mL 10%  $NaHCO_3$  sln. The organic layer was evaporated and purified by preparative tlc (1:60, MeOH:70%  $CH_2Cl_2$  / hexanes, 3 elutions) giving (+)-**36** (75.0 mg, 0.120 mmol, 55%) and (+)-**37** (34.0 mg, 0.055 mmol, 25%).

Indolocarbazole (+)-**36**. IR (thin film/NaCl) 3279.7 (brm), 3012.1 (m), 2952.1 (m), 2930.1 (m), 2850.1 (w), 1732.2 (m), 1646.2 (s), 1590.4 (m), 1513.7 (s), 1460.2 (s), 1260.3 (s), 1139.5 (s), 1028.1 (m), 744.5 (s)  $cm^{-1}$ ;  $^1H$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.26 (d,  $J=7.9$  Hz, 1H), 7.99 (d,  $J=7.7$  Hz, 1H), 7.92 (app.t,  $J=8.0$  Hz, 2H), 7.49 (app.t,  $J=7.7$  Hz, 1H), 7.47 (app.t,  $J=7.8$  Hz, 1H), 7.32 (app.t,  $J=7.9$  Hz, 1H), 7.30 (app.t,  $J=8.1$  Hz, 1H), 7.15 (dd,  $J=5.2, 6.9$  Hz, 1H), 7.02 (s, 1H), 6.94 (d,  $J=9.0$  Hz, 1H), 6.92 (d,  $J=9.0$  Hz, 1H), 6.35 (s, 1H), 5.0 (dd,  $J=17.8, 25.9$  Hz, 2H), 4.84 (dd,  $J=15.5, 17.5$  Hz, 2H), 3.92 (s, 3H), 3.74 (s, 3H), 3.71 (s, 3H), 3.39 (dd,  $J=7.3, 14.0$  Hz, 1H), 2.13 (s, 3H), 2.00 (dd,  $J=4.7, 14.0$  Hz, 1H);  $^{13}C$  NMR (62.5 MHz, DMSO- $d_6$ )  $\delta$  172.6, 168.6, 148.9, 148.2, 139.8, 136.7, 130.4, 130.0, 128.2, 125.3, 125.3, 124.8, 123.9, 123.8, 122.4, 120.9, 120.2, 119.8, 119.3, 118.9, 115.6, 114.6, 114.2, 112.3, 112.1, 108.8, 99.3, 84.8, 55.5, 52.4, 49.5, 45.4, 42.4, 22.6; HRMS (FAB)  $m/z$  calc'd for  $C_{36}H_{32}N_3O_7$  (M+H): 618.2240, found 618.2240;  $[\alpha]_D^{20}$  (+)-15° ( $c=0.1$ , MeOH).

Indolocarbazole (+)-**37**. IR (thin film/NaCl) 3462.3 (brm), 3014.0 (m), 2952.3 (m), 2925.1 (m), 2849.7 (m), 1730.8 (s), 1645.0 (m), 1514.7 (m), 1455.6 (s), 1403.9 (m), 1348.5 (m), 1312.6 (m), 1257.2 (s), 1235.0 (s), 1138.1 (s), 1068.8 (m), 1027.3 (m), 750.3 (s)  $cm^{-1}$ ;  $^1H$  NMR (500 MHz, DMSO- $d_6$ )  $\delta$  9.54 (d,  $J=7.9$  Hz, 1H), 8.01 (d,  $J=7.9$  Hz, 1H), 7.94 (d,  $J=8.2$  Hz, 1H), 7.89 (d,  $J=8.5$  Hz, 1H), 7.50 (app.t,  $J=7.5$  Hz, 1H), 7.45 (app.t,  $J=7.5$  Hz, 1H), 7.30 (app.t,  $J=7.5$  Hz, 1H), 7.29 (app.t,  $J=7.6$  Hz, 1H), 7.14 (dd,  $J=5.0, 7.2$  Hz, 1H), 7.01 (d,  $J=0.71$  Hz, 1H), 6.92 (app.t,  $J=8.2$  Hz, 1H), 6.92 (dd,  $J=1.1, 8.4$  Hz, 1H), 6.34 (bs, 1H), 4.97 (dd,  $J=17.9, 21.3$  Hz, 2H), 4.83 (dd,  $J=15.1, 21.9$  Hz, 2H), 3.92 (s, 3H), 3.74 (s, 3H), 3.71 (s, 3H), 3.40

(dd,  $J=7.5, 14.0$  Hz, 1H), 2.14 (s, 3H), 2.05 (dd,  $J=4.8, 14.0$  Hz, 1H);  $^{13}\text{C}$  NMR (62.5 MHz, DMSO- $d_6$ )  $\delta$  172.6, 168.9, 149.0, 148.2, 139.7, 136.8, 130.4, 126.2, 126.1, 125.4, 125.1, 124.9, 124.3, 122.0, 121.3, 120.2, 119.8, 119.2, 118.7, 116.3, 113.9, 113.8, 112.3, 112.1, 109.4, 99.3, 84.9, 84.8, 55.5, 52.4, 49.0, 45.4, 42.5, 22.8; HRMS (FAB)  $m/z$  calc'd for  $\text{C}_{36}\text{H}_{32}\text{N}_3\text{O}_7$  (M+H): 618.2240, found 618.2240;  $[\alpha]_D^{20}$  (+)-13° ( $c=0.1$ , MeOH).

(+)-K252a (14). To a stirred solution of (+)-36 (17.0 mg, 0.028 mmol) in  $\text{CH}_2\text{Cl}_2$  (1.4 mL) at 25 °C was added thioanisole (0.16 mL, 1.4 mmol, 50 eq) followed by 2,2,2-trifluoroacetic acid (1.4 mL). The solution was stirred for 6 h, at which point 2.0 mL sat.  $\text{NaHCO}_3$  sln. was added dropwise to neutralize the reaction mixture, the organic layer was separated, evaporated and purified via preparative tlc (1:40, MeOH:50%  $\text{CH}_2\text{Cl}_2$  / hexanes, 3 elutions) giving (+)-K252a (10.8 mg, 0.023 mmol, 83%).

Preparation of desamido K252a (40) To a suspension of indolo[2.3-a]carbazole (10) (1.0 g, 3.9 mmol) in 1,2-dichloroethane (130 mL) was added furanose 9 (1.8 g, 8.2 mmol) followed by CSA (100 mg, 0.43 mmol). The suspension was heated at reflux for 48 h, following which the reaction was allowed to cool to room temperature and was diluted with  $\text{CH}_2\text{Cl}_2$  (100mL), washed with 10%  $\text{NaHCO}_3$  (aq.), dried over  $\text{Na}_2\text{SO}_4$  and chromatographed on silica gel using 3:1 hexanes:ethyl acetate as eluent to afford indolocarbazole 40 (1.37 g, 85%) as a yellow foam.  $^1\text{H}$  NMR (500 MHz, acetone- $d_6$ ):  $\delta$  8.18 (app.t,  $J=6.6$  Hz, 1H), 8.18 (app.t,  $J=5.4$  Hz, 1H), 8.00 (m, 2H), 7.89 (d,  $J=8.5$  Hz, 1H), 7.75 (d,  $J=8.2$  Hz, 1H), 7.44 (td,  $J=0.9, 7.6$  Hz, 1H), 7.38 (td,  $J=1.0, 7.9$  Hz, 1H), 7.26 (app.t,  $J=6.9$  Hz, 1H), 7.25 (app.t,  $J=7.1$  Hz, 1H), 7.10 (dd,  $J=4.9, 7.3$  Hz, 1H), 5.18 (s, 1H), 3.99 (s, 3H), 3.44 (dd,  $J=7.5, 14.0$  Hz, 1H), 2.21 (s, 3H), 2.19 (dd,  $J=4.9, 14.0$  Hz, 1H).  $^{13}\text{C}$  NMR (125 MHz, acetone- $d_6$ ):  $\delta$  174.1, 140.8, 138.1, 127.7, 127.0, 125.6, 125.6, 125.5, 125.4, 121.6, 121.5, 121.2, 120.5, 120.4, 120.3, 115.0, 113.1, 112.8, 109.6, 99.9, 86.1, 86.0, 53.3, 43.2, 23.3. IR (thin film/NaCl): 3501.3 (brm), 3047.5 (m), 3006.7 (m), 2950.6 (m), 1729.4 (s), 1640.2 (m), 1568.1 (m), 1441.1 (s), 1305.9 (s), 1230.3 (s), 1128.1 (s), 740.0 (s)  $\text{cm}^{-1}$ . HRMS (EI)  $m/z$  Calc'd for  $\text{C}_{25}\text{H}_{20}\text{N}_2\text{O}_4$ : 412.1423. Found: 412.1419.

Preparation of aldehyde 41 To a stirred solution of ester 11 (1.0 g, 2.43 mmol) in THF (24.3mL), was added  $\text{LiBH}_4$  (106 mg, 4.85 mmol) at room

temperature. After 20 min the solvent was removed in vacuo. To the white residue was added 50 mL 1.0 N HCl on an ice bath. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x 50mL). The combined organic phases were dried with Na<sub>2</sub>SO<sub>4</sub> and chromatographed on silica gel using 1:1  
5 hexanes:ethyl acetate as eluent to afford a diol (815 mg, 87%) as a white solid. <sup>1</sup>H NMR (500 MHz, acetone-d<sub>6</sub>): δ 8.18 (d, J=7.6 Hz, 1H), 8.15 (d, J=7.8 Hz, 1H), 7.96 (s, 2H), 7.89 (d, J=8.5 Hz, 1H), 7.65 (d, J=8.1 Hz, 1H), 7.42 (app.t, J=7.6 Hz, 1H), 7.36 (app.t, J=8.2 Hz, 1H), 7.25 (app.t, J=7.6 Hz, 1H), 7.23 (app.t, J=7.4 Hz, 1H), 6.91 (dd, J=5.2, 7.4 Hz, 1H), 4.57 (s,  
10 1H), 4.18 (app.t, J=5.9 Hz, 1H), 4.06 (dd, J=5.4, 11.1 Hz, 1H), 3.90 (dd, J=7.1, 11.1 Hz, 1H), 3.30 (dd, J=7.6, 13.8 Hz, 1H), 2.23 (dd, J=5.1, 13.8 Hz, 1H), 2.22 (s, 3H). <sup>13</sup>C NMR (125 MHz, acetone-d<sub>6</sub>): δ 140.2, 137.4, 127.6, 126.3, 125.4, 125.0, 124.6, 124.6, 120.7, 120.6, 119.9, 119.5, 114.6, 112.2, 112.0, 108.8, 100.1, 84.2, 83.8, 65.5, 40.6, 21.5. IR (thin film/NaCl): 3416.8  
15 (brs), 3052.9 (m), 3010.5 (m), 2955.4 (w), 1732.7 (w), 1640.9 (m), 1568.5 (m), 1492.6 (m), 1459.0 (s), 1441.4 (s), 1309.0 (s), 1233.1 (s), 1031.9 (s), 741.0 (s) cm<sup>-1</sup>. HRMS (EI) *m/z* Calc'd for C<sub>24</sub>H<sub>20</sub>N<sub>2</sub>O<sub>3</sub>: 384.1474. Found: 384.1472. To a stirred solution of the diol (500 mg, 1.3 mmol) in 1:1 benzene:DMSO (8.7 mL) was added pyridinium trifluoroacetate (250 mg,  
20 1.3 mmol) followed by 1,3-dicyclohexylcarbodiimide (810 mg, 3.9 mmol). The flask was then quickly sealed with a septum, evacuated, and flushed with N<sub>2</sub> (3x). The heterogeneous mixture was stirred for 7h until reaction was complete as indicated by TLC. Benzene (15 mL) was added to the mixture and the 1,3-dicyclohexylurea (DCU) precipitate was filtered. The  
25 filtrate was washed with H<sub>2</sub>O (3x20 mL), and the combined aqueous layers were back extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x30mL). All organic layers were combined, dried with Na<sub>2</sub>SO<sub>4</sub>, and evaporated to an oily residue. A minimum amount of acetone (2 mL) was added to precipitate the remaining DCU. Filtration and evaporation to a yellow oil, which was purified by  
30 MPLC (3:1 hexanes:ethyl acetate) gave aldehyde 41 (375 mg, 73%, 63% 2 steps) as a yellow powder. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 9.70 (s, 1H), 7.99 (app.t, J=7.3 Hz, 2H), 7.78 (s, 2H), 8.02 (d, J=8.4 Hz, 1H), 7.29 (app.t, J=7.4 Hz, 1H), 7.24 (app.t, J=7.2 Hz, 1H), 7.22 (d, J=8.4 Hz, 1H), 7.17 (app.t, J=7.9 Hz, 1H), 7.15 (app.t, J=7.2 Hz, 1H), 6.59 (dd, J=5.0, 7.4 Hz,  
35 1H), 3.08 (s, 1H), 2.76 (dd, J=7.6, 14.6 Hz, 1H), 1.99 (s, 3H), 1.83 (dd, J=5.0, 14.7 Hz, 1H). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>): δ 199.4, 139.3, 136.9, 126.3, 126.3, 125.1, 124.7, 124.1, 121.2, 121.1, 120.8, 120.3, 120.3, 119.9, 113.1,

112.9, 112.2, 108.0, 97.7, 87.7, 84.0, 39.7, 23.0. IR (thin film/NaCl): 3486.7 (brn), 3054.6 (m), 3007.7 (m), 2945.3 (m), 2843.4 (w), 1723.9 (m), 1641.8 (m), 1568.6 (m), 1458.7 (m), 1441.1 (s), 1309.2 (s), 1232.5 (s), 1128.8 (m), 1004.2 (m), 741.7 (s)  $\text{cm}^{-1}$ . HRMS (EI)  $m/z$  Calc'd for  $\text{C}_{24}\text{H}_{18}\text{N}_2\text{O}_3$ : 382.1317. Found: 382.1319.

Preparation of hydroxy ketone 43 To a suspension of aldehyde 12 (75 mg, 0.196 mmol) in  $\text{Et}_2\text{O}$  (5.0 mL) was added  $\text{BF}_3 \cdot \text{OEt}_2$  (27  $\mu\text{L}$ , 0.216 mmol) and the mixture stirred vigorously for 6h.  $\text{CH}_2\text{Cl}_2$  (25 mL) was added to solubilize the suspension and the resulting solution was evaporated onto  $\text{SiO}_2$  (100mg) and chromatographed using 2:1 hexanes:ethyl acetate as eluent to provide ketone 43 (47 mg, 60%) as a white powder.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ ):  $\delta$  8.15 (d,  $J=7.7$  Hz, 1H), 8.10 (d,  $J=7.7$  Hz, 1H), 7.97 (d,  $J=8.5$  Hz, 1H), 7.92 (d,  $J=8.2$  Hz, 1H), 7.90 (d,  $J=8.2$  Hz, 1H), 7.43 (app.t,  $J=7.7$  Hz, 1H), 7.39 (app.t,  $J=7.8$  Hz, 1H), 7.32 (app.t,  $J=7.4$  Hz, 1H), 7.28 (app.t,  $J=7.5$  Hz, 1H), 7.25 (d,  $J=8.1$  Hz, 1H), 7.06 (d,  $J=7.3$  Hz, 1H), 4.89 (d,  $J=6.0$  Hz, 1H), 3.55 (dd,  $J=7.5$ , 14.3 Hz, 1H), 3.49 (d,  $J=6.5$  Hz, 1H), 2.99 (d,  $J=14.4$  Hz, 1H), 2.54 (s, 3H).  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ ):  $\delta$  199.8, 140.2, 136.3, 126.4, 125.7, 125.4, 125.1, 124.8, 124.6, 121.4, 120.8, 120.4, 120.2, 119.8, 115.2, 112.7, 112.4, 112.1, 107.9, 100.3, 84.03, 81.6, 44.7, 29.5. IR (thin film/NaCl): 3328.6 (brn), 3048.0 (w), 2923.7 (m), 2852.1 (w), 1731.4 (s), 1637.4 (m), 1441.5 (s), 1395.3 (m), 1312.0 (s), 1130.1 (m), 740.8 (s)  $\text{cm}^{-1}$ . HRMS (EI)  $m/z$  Calc'd for  $\text{C}_{24}\text{H}_{18}\text{N}_2\text{O}_3$ : 382.1317. Found: 382.1315.

Preparation of methoxy ketone 44 Montmorillonite Clay K-10 (1.2 g) was premixed with trimethylorthoformate (1.78 mL, 16.3 mmol) and immediately transfer to a stirred solution of aldehyde 12 (414 mg, 1.1 mmol) in  $\text{CHCl}_3$  (11 mL) aided by an additional 3 mL  $\text{CHCl}_3$ . The reaction was monitored by TLC (3:1 hexanes:ethyl acetate) and after approximately 0.5 h formation of the dimethyl acetal 16 was complete. The reaction mixture was filtered, and the filtrate evaporated *in vacuo*. The residue was dissolved in diethyl ether (110 mL) under an inert atmosphere, followed by addition of  $\text{BF}_3 \cdot \text{OEt}_2$  (2.85 mL, 23.1 mmol). The mixture was stirred for 4 d at 25  $^\circ\text{C}$ , following which triethyl amine (6.1 mL) and  $\text{CH}_2\text{Cl}_2$  (100 mL) were added, the solution was evaporated under reduced pressure and chromatographed on silica gel using 2:1 hexanes:ethyl acetate as eluent to afford methoxy ketone 44 (214 mg, 50%) as a yellow foam.  $^1\text{H}$  NMR (500

MHz, DMSO-d<sub>6</sub>):  $\delta$  8.21 (d, J=7.7 Hz, 1H), 8.16 (d, J=7.8 Hz, 1H), 7.97 (d, J=8.2 Hz, 1H), 7.95 (d, J=8.2 Hz, 1H), 7.88 (d, J=8.6 Hz, 1H), 7.68 (d, J=8.1 Hz, 1H), 7.46 (td, J=1.0, 7.4 Hz, 1H), 7.37 (td, J=1.1, 7.7 Hz, 1H), 7.36 (d, J=7.2 Hz, 1H), 7.30 (app.t, J=7.6 Hz, 1H), 7.23 (app.t, J=7.4 Hz, 1H), 5.02 (s, 1H), 3.94 (dd, J=7.2, 13.7 Hz, 1H), 3.39 (s, 3H), 2.62 (d, J=13.9 Hz, 1H), 2.52 (s, 3H). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>):  $\delta$  199.8, 139.4, 135.7, 125.0, 124.8, 124.5, 124.1, 124.0, 120.0, 119.8, 119.4, 119.2, 114.9, 112.1, 111.3, 109.2, 99.0, 88.2, 84.4, 58.9, 45.4, 29.2. IR (thin film/NaCl): 3046.6 (brm), 3003.8 (brw), 2927.9 (m), 2835.6 (m), 1736.6 (s), 1640.5 (m), 1565.8 (m), 1492.7 (m), 1442.9 (s), 1311.5 (s), 1144.3 (m), 1126.1 (s), 740.2 (s) cm<sup>-1</sup>. HRMS (EI) *m/z* Calc'd for C<sub>25</sub>H<sub>20</sub>N<sub>2</sub>O<sub>3</sub>: 396.1474. Found: 396.1474.

Preparation of desamido TAN-1030a (47) A suspension of ketone 15 (30 mg, 0.08 mmol), hydroxylamine hydrochloride (17 mg, 0.24 mmol), and NaOAc (20 mg, 0.24 mmol) in 50% aqueous EtOH (2.0 mL) was heated gently to reflux for 30 min. Following cooling to room temperature, solvent was removed *in vacuo*, and the residue purified by MPLC (2:1 hexanes:ethyl acetate) to provide oxime 47 (26 mg, 85%) as a yellow powder. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>):  $\delta$  10.43 (s, 1H), 8.17 (d, J=7.8 Hz, 1H), 8.13 (d, J=7.4 Hz, 1H), 7.91 (d, J=8.4 Hz, 1H), 7.89 (d, J=8.4 Hz, 1H), 7.88 (d, J=8.4 Hz, 1H), 7.67 (d, J=8.2 Hz, 1H), 7.44 (app.t, J=7.6 Hz, 1H), 7.34 (app.t, J=7.7 Hz, 1H), 7.27 (app.t, J=7.5 Hz, 1H), 7.20 (app.t, J=7.4 Hz, 1H), 6.98 (d, J=5.5 Hz, 1H), 4.70 (s, 1H), 3.61 (d, J=14.1 Hz, 1H), 3.42 (s, 3H), 2.97 (dd, J=5.7, 14.3 Hz, 1H), 2.42 (s, 3H). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>):  $\delta$  145.3, 139.3, 135.9, 126.0, 125.1, 124.9, 124.6, 124.2, 124.0, 120.0, 119.6, 119.4, 119.1, 119.1, 115.0, 111.8, 111.0, 109.1, 95.9, 83.7, 82.2, 58.3, 29.7, 28.4. IR (thin film/NaCl): 3249.5 (brm), 2918.3 (s), 2848.4 (s), 1728.1 (m), 1640.2 (m), 1443.1 (s), 1398.1 (m), 1312.0 (m), 1124.5 (s), 740.7 (s) cm<sup>-1</sup>. HRMS (EI) *m/z* Calc'd for C<sub>25</sub>H<sub>21</sub>N<sub>3</sub>O<sub>3</sub>: 411.1583. Found: 411.1582.

Preparation of desamido RK-286c (46) To a stirred solution of ketone 15 (12 mg, 0.03 mmol) in 1:1 MeOH: CH<sub>2</sub>Cl<sub>2</sub> (1.0 mL) was added NaBH<sub>4</sub> (3 mg, 0.08 mmol) at room temperature. After 5 minutes solvent was removed under reduced pressure. To the white residue was added 1 mL 1.0 N HCl on an ice bath. The mixture was stirred for 15 min at 25 °C and extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x 1mL). The combined organic phases were dried with Na<sub>2</sub>SO<sub>4</sub> and chromatographed on silica gel using 2:1 hexanes:ethyl



-31-

acetate as eluent to afford alcohol **46** (12 mg, 95%) as a white solid. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 8.14 (d, J=7.7 Hz, 1H), 8.11 (d, J=7.7 Hz, 1H), 7.90 (d, J=8.2 Hz, 1H), 7.85 (d, J=8.2 Hz, 1H), 7.81 (d, J=8.5 Hz, 1H), 7.39 (td, J=1.0, 8.1 Hz, 1H), 7.35 (ddd, J=1.4, 7.1, 8.4 Hz, 1H), 7.25 (m, 3H), 6.54 (d, J=5.6 Hz, 1H), 4.34 (m, 1H), 3.66 (d, J=3.0 Hz, 1H), 3.53 (s, 3H), 2.71 (dd, J=14.9 Hz, 1H), 2.45 (m, 1H), 2.30 (s, 3H), 1.66 (bs, 1H). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>): δ 139.6, 136.6, 128.3, 127.2, 126.5, 126.2, 124.8, 124.4, 123.9, 120.5, 120.3, 119.6, 119.3, 114.9, 112.1, 110.9, 107.6, 90.6, 83.1, 79.7, 60.5, 57.4, 33.7, 29.9. IR (thin film/NaCl): 3528.3 (brm), 3048.1 (m), 3000.2 (m), 2928.4 (m), 1643.7 (m), 1564.8 (m), 1493.3 (m), 1445.1 (s), 1344.4 (m), 1311.6 (s), 1231.2 (s), 1109.5 (brs) cm<sup>-1</sup>. HRMS (EI) *m/z* Calc'd for C<sub>25</sub>H<sub>22</sub>N<sub>2</sub>O<sub>3</sub>: 398.1630. Found: 398.1633.

Preparation of aldehyde (+)-**50** To a stirred solution of ester (+)-**36** (150 mg, 0.243 mmol) in THF (2.5 mL) was added LiBH<sub>4</sub> (12 mg, 0.535 mmol) at room temperature. After 20 min solvent was removed in vacuo. To the white residue, 10.0 mL 1.0 N HCl was added on an ice bath. The aqueous phase was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x 20 mL). The combined organic phases were dried with Na<sub>2</sub>SO<sub>4</sub> and chromatographed on silica gel using 1:1 hexanes:ethyl acetate as eluent to afford a diol (124 mg, 89%) as a white solid. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>): δ 9.25 (d, J=7.9 Hz, 1H), 7.97 (d, J=7.2 Hz, 1H), 7.96 (d, J=8.1 Hz, 1H), 7.78 (d, J=8.3 Hz, 1H), 7.48 (app.t, J=7.6 Hz, 1H), 7.43 (app.t, J=7.8 Hz, 1H), 7.29 (app.t, J=7.1 Hz, 1H), 7.28 (app.t, J=7.2 Hz, 1H), 7.02 (s, 1H), 7.96 (dd, J=5.2, 7.2 Hz, 1H), 6.94 (s, 2H), 5.33 (s, 1H), 5.06 (t, J=5.6 Hz, 1H), 5.02 (d, J=17.7 Hz, 1H), 4.95 (d, J=17.6 Hz, 1H), 4.85 (d, J=15.9 Hz, 1H), 4.85 (d, J=15.7 Hz, 1H), 3.85-3.81 (m, 2H), 3.75 (s, 3H), 3.72 (s, 3H), 3.14 (dd, J=7.6, 13.7 Hz, 1H), 2.15 (s, 3H), 1.94 (dd, J=4.8, 13.7 Hz, 1H). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>): δ 168.9, 148.9, 148.1, 140.0, 136.7, 130.5, 130.2, 128.7, 125.4, 125.3, 124.6, 124.3, 123.8, 122.4, 120.9, 120.0, 119.8, 119.2, 118.5, 115.2, 114.9, 114.0, 112.1, 111.8, 108.7, 100.2, 83.5, 64.7, 55.5, 55.5, 49.6, 45.4, 40.2, 40.1, 21.3. IR (thin film/NaCl): 3343.8 (brm), 3001.5 (w), 2950.7 (m), 2926.1 (m), 1647.4 (s), 1588.0 (m), 1514.4 (m), 1459.7 (s), 1422.2 (m), 1399.6 (m), 1312.4 (m), 1138.0 (s), 744.7 (s) cm<sup>-1</sup>. [α]<sub>D</sub><sup>25</sup> +112° (c=0.1, MeOH). To a stirred solution of the diol (395 mg, 0.67 mmol) in 1:1 benzene:DMSO (4.6 mL) was added pyridinium trifluoroacetate (130 mg, 0.67 mmol) followed by 1,3-dicyclohexylcarbodiimide (415 mg, 2.01 mmol). The flask was then quickly

- sealed with a septum, evacuated, and flushed with N<sub>2</sub> (3x). The heterogeneous mixture was stirred for 9h until reaction was complete as indicated by TLC. Benzene (5.0 mL) was added to the mixture and the 1,3-dicyclohexylurea (DCU) precipitate was filtered. The filtrate was washed with H<sub>2</sub>O (3x 5.0 mL), and the combined aqueous layers were back extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x 10.0 mL). All organic layers were combined, dried with Na<sub>2</sub>SO<sub>4</sub>, and evaporated to an oily residue. A minimum amount of acetone (2 mL) was added to precipitate the remaining DCU. Filtration and evaporation to a yellow oil, which was purified by MPLC (2:1→1:1 hexanes:ethyl acetate) gave aldehyde (+)-50 (280 mg, 71%, 63% 2 steps) as a yellow powder. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>): δ 10.07 (s, 1H), 9.31 (d, J=7.9 Hz, 1H), 8.02 (d, J=8.5 Hz, 1H), 7.99 (d, J=7.7 Hz, 1H), 7.87 (d, J=8.2 Hz, 1H), 7.50 (app.t, J=8.1 Hz, 1H), 7.47 (app.t, J=8.2 Hz, 1H), 7.32 (app.t, J=8.1 Hz, 1H), 7.17 (dd, J=7.2, 4.8 Hz, 1H), 7.04 (s, 1H), 6.94 (d, J=9.6 Hz, 1H), 6.93 (d, J=8.1 Hz, 1H), 6.57 (bs, 1H), 5.02 (d, J=17.6 Hz, 1H), 4.98 (d, J=17.7 Hz, 1H), 4.87 (d, J=15.2 Hz, 1H), 4.83 (d, J=15.2 Hz, 1H), 3.76 (s, 3H), 3.73 (s, 3H), 3.24 (dd, J=7.6, 14.0 Hz, 1H), 2.22 (s, 3H), 2.00 (dd, J=4.5, 14.0 Hz, 1H). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>): δ 202.2, 168.7, 148.9, 148.1, 139.9, 136.9, 130.4, 130.2, 128.2, 125.5, 125.1, 123.9, 123.9, 122.5, 121.1, 120.4, 119.9, 119.6, 119.1, 115.8, 114.6, 114.4, 112.1, 111.8, 109.0, 98.7, 86.8, 84.3, 55.5, 55.5, 49.6, 45.5, 39.4, 22.7. IR (thin film/NaCl): 3253.9 (brm), 3010.7 (m), 2953.6 (m), 2934.0 (m), 2833.9 (s), 1734.0 (s), 1646.2 (s), 1614.7 (w), 1589.9 (m), 1514.1 (m), 1399.1 (s), 1275.7 (m), 1138.4 (s), 1024.8 (m), 745.1 (s) cm<sup>-1</sup>. [α]<sub>D</sub><sup>25</sup> +48° (c=0.1, MeOH).
- 25 Preparation of hydroxy ketone (+)-51 To a suspension of aldehyde (+)-50 (100 mg, 0.170 mmol) in Et<sub>2</sub>O (17.0 mL) was added BF<sub>3</sub>•OEt<sub>2</sub> (23 μL, 0.187 mmol) and the mixture stirred vigorously for 12h at 25-30 °C, when again was treated with BF<sub>3</sub>•OEt<sub>2</sub> (23 μL, 0.187 mmol) and stirred for an additional 12 h at the same temperature. The reaction mixture was filtered to provide ketone (+)-51 (85 mg, 85%) as a white powder. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>, 310 K): δ 9.35 (d, J=7.9 Hz, 1H), 8.06 (d, J=8.6 Hz, 1H), 7.92 (d, J=7.7 Hz, 1H), 7.72 (d, J=8.2 Hz, 1H), 7.53 (app.t, J=7.6 Hz, 1H), 7.43 (app.t, J=8.1 Hz, 1H), 7.40 (d, J=6.6 Hz, 1H), 7.35 (app.t, J=7.5 Hz, 1H), 7.29 (app.t, J=7.4 Hz, 1H), 7.02 (s, 1H), 6.93 (s, 2H), 6.12 (d, J=5.1 Hz, 1H), 5.23 (d, J=4.5 Hz, 1H), 4.96 (s, 2H), 4.85 (d, J=15.1 Hz, 1H), 4.81 (d, J=15.1 Hz, 1H), 3.97 (dd, J=6.7, 14.1 Hz, 1H), 3.75 (s, 3H), 3.72 (s, 3H),

2.66 (d,  $J=14.1$  Hz, 1H), 2.54 (s, 3H).  $^{13}\text{C}$  NMR (500 MHz, DMSO- $d_6$ ):  $\delta$  201.1, 168.6, 148.9, 148.1, 140.3, 136.0, 130.4, 129.8, 126.9, 125.6, 125.5, 124.9, 124.0, 123.6, 122.8, 120.7, 120.4, 119.9, 119.9, 118.8, 115.9, 115.1, 114.3, 112.1, 111.8, 109.2, 100.5, 84.4, 80.0, 55.5, 55.5, 49.6, 45.4, 44.9, 29.4. IR (thin film/NaCl): 3300.0 (brs), 2999.5 (brm), 2848.6 (m), 1728.9 (m), 1665.5 (s), 1503.3 (m), 1451.2 (s), 1406.8 (m), 1132.8 (s), 1021.9 (m), 750.6 (s)  $\text{cm}^{-1}$ .  $[\alpha]_{\text{D}}^{25} +83^\circ$  ( $c=0.1$ , DMSO).

Preparation of Ester (+)-35 To a solution of ketone (+)-51 (10 mg, 0.017 mmol) in 1:1 MeOH/ $\text{CH}_2\text{Cl}_2$  (1.0 mL) was added Copper (I) chloride (30 mg, 0.30 mmol), and the mixture warmed to reflux for 15 min. Solvent was removed *in vacuo* and the resulting residue subjected to silica gel chromatography (1:1, hexanes:ethyl acetate) to afford (+)-36 (10 mg, 95%) as a colorless solid.

Preparation of diol (+)-53 To a stirred solution of ketone (+)-51 (85 mg, 0.15 mmol) in 1:1:2 MeOH: $\text{CH}_2\text{Cl}_2$ : $\text{CHCl}_3$  (20.0 mL),  $\text{NaBH}_4$  (20 mg, 0.53 mmol) was added at room temperature. After 5 minutes solvent was removed under reduced pressure. To the white residue, 10 mL 1.0 N HCl was added on an ice bath. The mixture was stirred for 15 min at 25  $^\circ\text{C}$  and extracted with  $\text{CH}_2\text{Cl}_2$  (3x 20 mL). The combined organic phases were dried with  $\text{Na}_2\text{SO}_4$  and chromatographed on silica gel using 1:1 hexanes:ethyl acetate as eluent to afford alcohol (+)-53 (81 mg, 95%) as a white solid.  $^1\text{H}$  NMR (500 MHz, acetone- $d_6$ ):  $\delta$  9.53 (d,  $J=7.9$  Hz, 1H), 8.11 (d,  $J=8.5$  Hz, 1H), 7.88 (d,  $J=7.7$  Hz, 1H), 7.51 (d,  $J=8.2$  Hz, 1H), 7.46 (app.t,  $J=7.2$  Hz, 1H), 7.36 (app.t,  $J=7.9$  Hz, 1H), 7.29 (app.t,  $J=7.4$  Hz, 1H), 7.22 (app.t,  $J=7.4$  Hz, 1H), 7.08 (s, 1H), 6.98 (d,  $J=8.3$  Hz, 1H), 6.91 (d,  $J=8.2$  Hz, 1H), 6.76 (d,  $J=5.1$  Hz, 1H), 4.95 (d,  $J=17.1$  Hz, 1H), 4.90 (d,  $J=71.1$  Hz, 1H), 4.89 (d,  $J=15.2$  Hz, 1H), 4.85 (d,  $J=15.2$  Hz, 1H), 4.24 (d,  $J=8.5$  Hz, 1H), 4.23 (bs, 1H), 4.14 (d,  $J=8.6$  Hz, 1H), 3.77 (s, 3H), 3.76 (s, 3H), 3.64 (bs, 1H), 2.76 (d,  $J=15.1$  Hz, 1H), 2.65 (d,  $J=15.1$  Hz, 1H), 2.35 (s, 3H).  $^{13}\text{C}$  NMR (125 MHz, acetone- $d_6$ ):  $\delta$  170.4, 150.6, 149.7, 141.2, 137.7, 132.0, 130.7, 130.4, 127.6, 127.1, 125.8, 125.3, 125.0, 124.3, 121.5, 121.0, 120.6, 120.0, 119.8, 116.6, 116.0, 115.0, 112.8, 108.9, 93.3, 80.6, 74.7, 65.4, 56.1, 50.4, 46.6, 35.4, 30.4. IR (thin film/NaCl): 3355.5 (brm), 2922.9 (m), 2847.8 (m), 1654.5 (s), 1501.5 (w), 1449.3 (s), 1254.5 (s), 1136.8 (s), 1025.7 (m), 747.1 (s)  $\text{cm}^{-1}$ .  $[\alpha]_{\text{D}}^{25} +37^\circ$  ( $c=0.1$ , MeOH).

- Preparation of methyl ether (+)-54 To a stirred suspension of NaH (14 mg, 0.58 mmol) in THF (1.0 mL) was added a solution of alcohol (+)-53 (81 mg, 0.138 mmol) in THF (7 mL). The resulting mixture was stirred for 10 min with the visible evolution of gas, and for an additional 15 min thereafter.
- 5 Addition of MeI (9.5  $\mu$ L, 0.15 mmol) produced a single product by TLC (2.5:1 hexanes:acetone). After approximately 50 min the reaction was quenched by addition of 1.0 mL 1.0N HCl followed by 2.0 mL H<sub>2</sub>O. Extraction of the solution with CH<sub>2</sub>Cl<sub>2</sub> (3x 10 mL), drying over Na<sub>2</sub>SO<sub>4</sub> and evaporation to a residue which could be purified by MPLC (2.5:1 hexanes:acetone) provided
- 10 methyl ether (+)-54 (67 mg, 80%) as a yellow foam. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>):  $\delta$  9.54 (d, J=7.9 Hz, 1H), 7.90 (d, J=8.5 Hz, 1H), 7.81 (d, J=7.7 Hz, 1H), 7.48 (app.t, J=7.6 Hz, 1H), 7.41 (app.t, J=7.2 Hz, 1H), 7.38 (app.t, J=7.2 Hz, 1H), 7.28 (m, 2H), 6.97 (d, J=8.2 Hz, 1H), 6.95 (s, 1H), 6.86 (d, J=8.1 Hz, 1H), 6.60 (d, J=5.8 Hz, 1H), 4.96 (d, J=15.0 Hz, 1H), 4.89 (d,
- 15 J=15.0 Hz, 1H), 4.84 (d, J=16.7 Hz, 1H), 4.79 (d, J=16.6 Hz, 1H), 4.38 (d, J=2.6 Hz, 1H), 3.88 (s, 3H), 3.85 (s, 3H), 3.71 (d, J=2.6 Hz, 1H), 3.57 (s, 3H), 2.76 (dd, J=3.1, 15.1 Hz, 1H), 2.50 (bd, J=14.7 Hz, 1H), 2.3 (s, 3H). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>, 315 K):  $\delta$  170.3, 149.6, 148.7, 140.1, 136.8, 130.8, 129.4, 127.0, 126.4, 125.3, 124.8, 124.3, 123.7, 120.7, 120.4, 120.2,
- 20 120.0, 119.6, 116.0, 115.5, 114.5, 111.6, 111.5, 107.1, 90.7, 83.2, 79.5, 60.6, 57.4, 56.1, 56.0, 49.9, 46.5, 33.6, 30.1. IR (thin film/NaCl): 3423.7 (brm), 2923.2 (s), 2848.1 (m), 2636.2 (m), 1647.2 (s), 1514.3 (m), 1462.9 (s), 1258.0 (m), 1235.3 (m), 1136.9 (m), 1026.9 (w), 743.3 (s) cm<sup>-1</sup>. [ $\alpha$ ]<sub>D</sub><sup>25</sup> +48° (c=0.1, MeOH).
- 25 Preparation of (+)-RK-286c (50) To a stirred solution of ether (+)-54 (10 mg, 0.017 mmol) in anisole or thioanisole (80  $\mu$ L,  $\approx$ 50 equiv) was added TFA (0.5 mL). The reaction was monitored by TLC, and after 24 h had proceeded to completion, whereupon 1.0 mL H<sub>2</sub>O was added, followed by extraction with CH<sub>2</sub>Cl<sub>2</sub> (3x 5mL). Combined organic layers were washed
- 30 with saturated aqueous NaHCO<sub>3</sub> (5 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated to a residue, which was purified by preparative TLC (5% MeOH:CH<sub>2</sub>Cl<sub>2</sub>) to provide (+)-RK-286c (50, 6 mg, 75%). <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>):  $\delta$  9.27 (d, J=7.9 Hz, 1H), 8.47 (bs, 1H), 7.99 (d, J=8.5 Hz, 1H), 7.94 (d, J=7.7 Hz, 1H), 7.59 (d, J=8.2 Hz, 1H), 7.45 (app.t, J=7.4 Hz,
- 35 1H), 7.40 (app.t, J=7.5 Hz, 1H), 7.26 (app.t, J=7.5 Hz, 1H), 6.78 (d, J=5.3

- Hz, 1H), 4.95 (d, J=17.6 Hz, 1H), 4.89 (d, J=17.7 Hz, 1H), 4.25 (bs, 1H), 4.17 (bs, 1H), 3.83 (d, J=2.7 Hz, 1H), 3.41 (s, 3H), 2.60 (ddd, J=3.2, 5.6, 14.8 Hz, 1H), 2.41 (dd, J=3.3, 14.8 Hz, 1H), 2.31 (s, 3H). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>): δ 139.7, 136.1, 129.5, 125.5, 124.7, 124.1, 123.9, 122.6, 120.6, 119.5, 118.9, 118.6, 115.7, 108.6, 90.9, 82.3, 79.5, 58.8, 56.4, 45.3, 33.9, 29.9. IR (thin film/NaCl): 3354.0 (brm), 2920.4 (s), 2851.6 (m), 1677.2 (s), 1636.0 (m), 1585.3 (m), 1456.2 (s), 1352.8 (s), 1318.7 (s), 1231.7 (m), 1117.3 (m), 743.8 (s) cm<sup>-1</sup>. [α]<sub>D</sub><sup>25</sup> +41.1° (c=0.18, EtOAc); natural RK-286c [α]<sub>D</sub><sup>20</sup> +45.3° (c=0.22, EtOAc).
- 10 Preparation of Diol (+)-55 To a stirred solution of ether (+)-54 (112 mg, 0.186 mmol) in CDCl<sub>3</sub> (2.0 mL) was added Martin's sulfurane (187 mg, 0.28 mmol). The reaction rapidly proceeded to a less polar product as monitored by TLC, and after 20 min was complete. Solvent was evaporated and the residue subjected to silica gel chromatography (2:1 hexanes:ethyl acetate) to provide an olefin (96 mg, 88%) as a white solid. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>, 315 K): δ 9.31 (d, J=7.9 Hz, 1H), 8.11 (d, J=8.6 Hz, 1H), 7.91 (d, J=7.7 Hz, 1H), 7.86 (d, J=8.2 Hz, 1H), 7.50 (td, J=1.0, 7.34 Hz, 1H), 7.43 (app.t, J=7.8 Hz, 1H), 7.31 (app.t, J=7.0 Hz, 1H), 7.28 (app.t, J=7.1 Hz, 1H), 7.13 (d, J=1.9 Hz, 1H), 7.02 (s, 1H), 6.93 (d, J=8.6 Hz, 1H), 6.92 (d, J=8.6 Hz, 1H), 6.09 (d, J=10.4 Hz, 1H), 5.77 (dt, J=2.3, 10.4 Hz, 1H), 4.95 (s, 2H), 4.85 (d, J=15.1 Hz, 1H), 4.81 (d, J=15.1 Hz, 1H), 4.48 (d, J=1.4 Hz, 1H), 3.74 (s, 3H), 3.71 (s, 3H), 3.57 (s, 3H), 2.20 (s, 3H). <sup>13</sup>C NMR (125 MHz, acetone-d<sub>6</sub>): δ 169.9, 150.5, 149.7, 141.3, 137.4, 131.8, 131.2, 130.5, 127.7, 127.1, 126.4, 126.2, 125.5, 125.3, 124.3, 121.5, 121.2, 121.1, 120.5, 120.4, 118.0, 117.1, 115.9, 112.8, 112.8, 109.1, 91.5, 80.8, 78.8, 57.7, 56.0, 56.0, 50.5, 46.5, 28.0. IR (thin film/NaCl): 2920.5 (s), 2851.5 (s), 1709.8 (m), 1674.3 (s), 1589.0 (m), 1513.7 (m), 1457.5 (s), 1222.9 (m), 1026.6 (m), 745.3 (m) cm<sup>-1</sup>. [α]<sub>D</sub><sup>25</sup> +36° (c=0.1, MeOH). To a stirred solution of 4-methylmorpholine N-oxide (6 mg, 0.05 mmol) and OsO<sub>4</sub> (0.6 mL of a 2.5% solution in *t*-BuOH, 0.05 mmol) in 4:1 acetone:H<sub>2</sub>O (2 mL) was added solution of the olefin (25 mg, 0.043 mmol) in acetone (1 mL). The reaction was monitored by TLC, and after 16 h had proceeded to completion, whereupon 100 mg NaHSO<sub>3</sub> was added in 1.0 mL H<sub>2</sub>O, and the black solution was stirred for 20 min and filtered, followed by extraction with CH<sub>2</sub>Cl<sub>2</sub> (3x 15mL). Combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated to a residue, which was purified by MPLC (1:1 hexanes:ethyl

acetate) to provide diol (+)-55 (23 mg, 84%) as a white powder. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>): δ 9.36 (d, J=7.9 Hz, 1H), 7.95 (d, J=8.6 Hz, 1H), 7.94 (d, J=7.6 Hz, 1H), 7.64 (d, J=8.1 Hz, 1H), 7.55 (app.t, J=7.6 Hz, 1H), 7.45 (app.t, J=7.7 Hz, 1H), 7.35 (app.t, J=7.5 Hz, 1H), 7.29 (app.t, J=7.5 Hz, 1H), 7.02 (s, 1H), 6.94 (s, 2H), 6.59 (d, J=1.6 Hz, 1H), 6.13 (d, J=3.8 Hz, 1H), 5.07 (d, J=6.0 Hz, 1H), 4.99 (d, J=17.8 Hz, 1H), 4.95 (d, J=17.8 Hz, 1H), 4.83 (s, 2H), 4.12 (d, J=10.1 Hz, 1H), 4.12 (dd, J=2.3, 3.8 Hz, 1H), 3.74 (s, 3H), 3.72 (s, 3H), 3.62 (s, 3H), 3.55 (ddd, J=2.3, 6.1, 10.1 Hz, 1H), 2.37 (s, 3H). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>): δ 168.8, 148.9, 148.1, 140.3, 136.5, 130.4, 129.9, 127.8, 125.7, 125.0, 124.7, 123.5, 122.7, 120.8, 120.2, 119.9, 119.9, 118.7, 115.5, 114.8, 114.1, 112.0, 111.7, 108.8, 95.6, 87.3, 83.1, 71.7, 65.6, 61.6, 55.5, 55.5, 49.6, 45.5, 29.0. IR (thin film/NaCl): 3411.2 (brm), 2929.3 (m), 2849.4 (w), 2656.3 (m), 1590.0 (m), 1514.0 (m), 1461.2 (s), 1350.9 (m), 1273.6 (s), 1127.1 (s), 1025.0 (m), 743.3 (s) cm<sup>-1</sup>. [α]<sub>D</sub><sup>25</sup> +17° (c=0.1, MeOH).

Preparation of (+)-MLR-52 (51) To a stirred solution of diol (+)-55 (10 mg, 0.016 mmol) in anisole or thioanisole (80 μL, ~50 equiv) was added TFA (0.5 mL). The reaction was monitored by TLC, and after 16 h had proceeded to completion, whereupon 1.0 mL H<sub>2</sub>O was added, followed by extraction with CH<sub>2</sub>Cl<sub>2</sub> (3x 5mL). Combined organic layers were washed with saturated aqueous NaHCO<sub>3</sub> (5 mL), dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated to a residue, which was purified by preparative TLC (5% MeOH:CH<sub>2</sub>Cl<sub>2</sub>) to provide (+)-MLR-52 (51, 6 mg, 77%). <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>): δ 9.31 (d, J=7.9 Hz, 1H), 8.61 (bs, 1H), 7.99 (d, J=7.7 Hz, 1H), 7.96 (d, J=8.7 Hz, 1H), 7.62 (d, J=8.2 Hz, 1H), 7.53 (app.t, J=7.5 Hz, 1H), 7.45 (td, J=0.8, 7.7 Hz, 1H), 7.32 (app.t, J=7.4 Hz, 1H), 7.32 (app.t, J=7.4 Hz, 1H), 6.58 (d, J=1.6 Hz, 1H), 6.12 (d, J=4.0 Hz, 1H), 5.06 (d, J=5.9 Hz, 1H), 4.99 (d, J=17.6 Hz, 1H), 4.95 (d, J=17.5 Hz, 1H), 4.13 (d, J=10.3 Hz, 1H), 4.12 (dd, J=1.6, 2.6 Hz, 1H), 3.62 (s, 3H), 3.56 (ddd, J=2.6, 6.2, 10.3 Hz, 1H). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>): δ 171.8, 140.2, 136.4, 132.6, 127.8, 125.8, 125.5, 124.8, 124.6, 123.6, 122.7, 120.9, 120.1, 119.7, 119.3, 115.4, 114.9, 114.3, 108.7, 95.6, 87.2, 83.1, 71.7, 65.6, 61.6, 45.4, 29.0. IR (thin film/NaCl): 3348.5 (brm), 2922.9 (s), 2851.9 (m), 1638.2 (s), 1586.6 (m), 1455.5 (s), 1373.5 (m), 1336.6 (m), 1320.8 (m), 1275.0 (m), 1224.7 (m), 1200.3 (w), 1119.5 (s), 740.8 (s,) cm<sup>-1</sup>. [α]<sub>D</sub><sup>25</sup> +65° (c=0.1, MeOH); natural MLR-52 [α]<sub>D</sub> +68° (c=0.093, MeOH).

Preparation of oxime (-)-56 A suspension of ketone (+)-51 (100 mg, 0.17 mmol), hydroxylamine hydrochloride (165 mg, 2.38 mmol), and NaOAc (167 mg, 2.04 mmol) in 80% aqueous EtOH (35.0 mL) was heated gently to reflux for 30 min. Following cooling to room temperature, solvent was removed *in vacuo*, and the residue purified by MPLC (1:1 hexanes:ethyl acetate) to provide oxime (-)-56 (98 mg, 95%) as a yellow powder. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>): δ 10.3 (s, 1H), 9.34 (d, J=7.9 Hz, 1H), 8.08 (d, J=8.6 Hz, 1H), 7.90 (d, J=7.6 Hz, 1H), 7.71 (d, J=8.3 Hz, 1H), 7.51 (app.t, J=7.6 Hz, 1H), 7.42 (app.t, J=7.9 Hz, 1H), 7.32 (app.t, J=7.7 Hz, 1H), 7.28 (app.t, J=7.4 Hz, 1H), 7.04 (d, J=6.3 Hz, 1H), 7.03 (s, 1H), 6.95 (d, J=8.4 Hz, 1H), 6.93 (d, J=8.2 Hz, 1H), 5.56 (m, 2H), 4.97 (d, J=18.1 Hz, 1H), 4.93 (d, J=16.9 Hz, 1H), 4.85 (d, J=15.0 Hz, 1H), 4.45 (d, J=15.0 Hz, 1H), 3.75 (s, 3H), 3.72 (s, 3H), 3.61 (d, J=13.9 Hz, 1H), 3.01 (dd, J=5.8, 14.3 Hz, 1H), 2.46 (s, 3H). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>): δ 168.8, 148.9, 148.1, 147.4, 140.2, 136.1, 130.5, 129.6, 128.1, 125.4, 125.3, 124.7, 124.6, 123.6, 122.8, 120.5, 120.1, 119.9, 119.6, 118.5, 116.0, 114.8, 113.9, 112.1, 111.9, 108.9, 97.4, 82.0, 74.9, 55.5, 55.5, 49.5, 45.5, 29.6, 28.6. IR (thin film/NaCl): 3324.0 (brm), 2995.0 (w), 2911.3 (m), 1660.0 (s), 1589.7 (m), 1513.5 (s), 1461.1 (s), 1417.9 (m), 1399.0 (m), 1349.2 (s), 1315.5 (m), 1260.0 (s), 1234.6 (m), 1124.4 (m), 1027.2 (m), 741.7 (s) cm<sup>-1</sup>. [α]<sub>D</sub><sup>20</sup> -18° (c=0.1, CH<sub>2</sub>Cl<sub>2</sub>).

Preparation of methoxy oxime (-)-57 To a mixture of oxime (-)-56 (90 mg, 0.15 mmol), MeI (88 μL, 1.42 mmol), and powdered KOH (88 mg, 1.58 mmol) was added *n*-Bu<sub>4</sub>NBr (10 mg, 0.03 mmol). The mixture was stirred under N<sub>2</sub> for 30 min, solvent was removed *in vacuo*, and the residue was subjected to silica gel chromatography (1:1 hexanes:ethyl acetate) to provide methoxime (-)-57 (85 mg, 90%) as a yellow powder. <sup>1</sup>H NMR (500 MHz, DMSO-d<sub>6</sub>, 345 K): δ 9.36 (d, J=8.0 Hz, 1H), 7.99 (d, J=8.6 Hz, 1H), 7.93 (d, J=7.8 Hz, 1H), 7.68 (d, J=8.3 Hz, 1H), 7.51 (app.t, J=7.6 Hz, 1H), 7.44 (app.t, J=7.8 Hz, 1H), 7.33 (app.t, J=7.2 Hz, 1H), 7.30 (app.t, J=7.1 Hz, 1H), 7.04 (s, 1H), 7.02 (d, J=5.6 Hz, 1H), 6.97 (d, J=9.4 Hz, 1H), 6.94 (d, J=8.1 Hz, 1H), 4.97 (s, 2H), 4.86 (d, J=15.5 Hz, 1H), 4.85 (d, J=15.7 Hz, 1H), 4.76 (s, 1H), 3.76 (s, 3H), 3.74 (s, 3H), 3.54 (d, J=14.4 Hz, 1H), 3.45 (s, 3H), 3.16 (dd, J=5.9, 14.4 Hz, 1H), 3.14 (s, 3H), 2.46 (s, 3H). <sup>13</sup>C NMR (125 MHz, DMSO-d<sub>6</sub>): δ 168.7, 148.9, 148.1, 147.3, 139.8, 136.1, 130.4,

129.5, 128.0, 125.4, 125.3, 124.7, 124.6, 123.6, 122.7, 120.6, 120.2, 119.9, 119.6, 118.6, 115.5, 114.9, 113.8, 112.2, 112.0, 108.9, 96.1, 83.3, 82.0, 60.8, 58.4, 55.5, 55.5, 49.5, 45.4, 30.4, 28.5. IR (thin film/NaCl): 2998.0 (w), 2926.3 (m), 1674.1 (s), 1590.0 (m), 1513.7 (s), 1460.9 (s), 1418.2 (m), 1397.9 (s), 1349.4 (s), 1316.2 (s), 1262.1 (m), 1225.6 (m), 1044.3 (m), 743.5 (m)  $\text{cm}^{-1}$ .  $[\alpha]_D^{25} -22^\circ$  ( $c=0.1$ ,  $\text{CH}_2\text{Cl}_2$ ).

Preparation of amine (+)-58 A mixture of oxime (+)-57 (85 mg, 0.13 mmol) and  $\text{PtO}_2$  (28 mg) in a 60% aqueous acetic acid (15.0 mL) was treated with  $\text{H}_2$ , and the reaction was monitored by TLC (1:1 hexanes:ethyl acetate). Upon completion, the mixture was filtered through celite and the filtrate was evaporated. The residue was dissolved in  $\text{CH}_2\text{Cl}_2$  (40 mL) and washed with 8.0 mL 1.0N NaOH. The aqueous layer was then back-extracted with  $\text{CH}_2\text{Cl}_2$  (2x 15 mL). The combined organic layers were dried over  $\text{Na}_2\text{SO}_4$  and evaporated to a residue (79 mg), which was used in the next step without further purification. An analytical sample of primary amine (+)-58 could be obtained by preparative TLC of the above residue using 5%  $\text{MeOH}:\text{CH}_2\text{Cl}_2$  as eluent.  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ , 310 K):  $\delta$  9.55 (d,  $J=7.9$  Hz, 1H), 7.95 (d,  $J=8.5$  Hz, 1H), 7.83 (d,  $J=7.7$  Hz, 1H), 7.51 (app.t,  $J=7.6$  Hz, 1H), 7.42 (app.t,  $J=8.2$  Hz, 1H), 7.40 (app.t,  $J=7.5$  Hz, 1H), 7.30 (app.t,  $J=7.8$  Hz, 2H), 6.99 (d,  $J=9.4$  Hz, 2H), 6.87 (d,  $J=8.0$  Hz, 1H), 6.59 (d,  $J=4.9$  Hz, 1H), 4.98 (d,  $J=14.9$  Hz, 1H), 4.92 (d,  $J=14.9$  Hz, 1H), 4.87 (d,  $J=16.7$  Hz, 1H), 4.82 (d,  $J=16.7$  Hz, 1H), 3.88 (s, 3H), 3.86 (s, 3H), 3.75 (m, 2H), 3.46 (s, 3H), 2.63 (m, 2H), 2.32 (s, 3H), 1.27 (bs, 2H).  $^{13}\text{C}$  NMR (125 MHz,  $\text{CDCl}_3$ , 315 K):  $\delta$  170.2, 149.6, 148.7, 140.1, 137.0, 130.8, 129.6, 129.5, 127.0, 126.2, 125.4, 124.7, 124.5, 123.8, 120.8, 120.5, 120.2, 120.2, 119.6, 116.0, 115.4, 114.6, 111.6, 111.6, 107.4, 91.3, 84.2, 80.2, 57.5, 56.1, 56.1, 49.9, 46.5, 42.6, 34.6, 30.0. IR (thin film/NaCl): 3414.7 (brw), 2920.8 (s), 2851.7 (s), 1733.7 (w), 1672.8 (s), 1636.0 (w), 1588.1 (m), 1513.5 (s), 1352.7 (s), 1259.3 (s), 1136.7 (m), 744.2 (m)  $\text{cm}^{-1}$ .  $[\alpha]_D^{25} +14.3^\circ$  ( $c=0.14$ ,  $\text{CHCl}_3$ ).

Preparation of (+)-staurosporine (49) Crude amine (+)-58 was dissolved in THF (2.0 mL) and treated with formic acetic anhydride in THF (1.3  $\mu\text{L}$  of a 1.3 M solution in THF, 0.17 mmol)(FAA prepared by treatment of 1.0 equiv acetic anhydride with 1.2 equiv formic acid followed by reflux for 2 h). TLC analysis showed rapid formation of a less polar substance. A stream of  $\text{N}_2$



was used to evaporate the solvent, followed by high vacuum for 15 min. THF (1.3 mL) was added to dissolve the residue, the reaction vessel was cooled to 0 °C, and BH<sub>3</sub>•DMS (193 µL of a 2.0 N solution in toluene, 0.39 mmol) was introduced. The solution was heated to reflux for 2 h at which point it was again cooled to 0 °C. Methanolic HCl (1.0 mL) was added along with excess MeOH (1.3 mL) and the solution was refluxed for an additional hour. Following cooling, volatiles were removed *in vacuo*, and the solid residue was azeotroped with MeOH (5x 5.0 mL). To the remaining residue was added 7.0 mL CH<sub>2</sub>Cl<sub>2</sub> followed by 1.0 N NaOH (5.0 mL), layers were separated, and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3x 7.0 mL). Combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, evaporated, and purified by MPLC (5% MeOH:CH<sub>2</sub>Cl<sub>2</sub>) to give a methyl amine (80 mg, 91% 2 steps from 14) as a yellow foam. <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>, 320 °K): δ 9.55 (d, J=7.9 Hz, 1H), 7.89 (d, J=8.5 Hz, 1H), 7.82 (d, J=7.3 Hz, 1H), 7.48 (td, J=1.0, 7.5 Hz, 1H), 7.39 (td, J=1.0, 7.4 Hz, 1H), 7.38 (app.t, J=7.3 Hz, 1H), 7.27 (m, 2H), 7.01 (m, 2H), 6.88 (d, J=8.7 Hz, 1H), 6.57 (dd, J=1.4, 6.0 Hz, 1H), 4.98 (d, J=14.9 Hz, 1H), 4.91 (d, J=14.9 Hz, 1H), 4.84 (s, 2H), 3.92 (d, J=3.0 Hz, 1H), 3.88 (s, 3H), 3.87 (s, 3H), 3.37 (dd, J=3.8, 7.7 Hz, 1H), 3.33 (bs, 3H), 2.72 (ddd, J=1.3, 4.6, 14.5 Hz, 1H), 2.46 (m, 1H), 2.35 (s, 3H). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>): δ 170.4, 149.3, 148.4, 139.6, 136.7, 130.6, 130.4, 129.3, 127.1, 126.6, 125.1, 124.5, 124.3, 123.5, 120.7, 120.4, 120.0, 119.8, 119.1, 115.5, 114.9, 114.0, 111.2, 111.2, 107.0, 91.2, 83.9, 80.2, 57.5, 56.0, 55.9, 50.7, 49.9, 46.4, 33.2, 30.1, 29.9. IR (thin film/NaCl): 2954.1 (m), 2915.1 (m), 1673.2 (s), 1635.8 (m), 1462.7 (s), 1399.0 (s), 1352.6 (s), 1258.7 (m), 1136.5 (m), 1026.9 (m), 745.2 (s) cm<sup>-1</sup>. [α]<sub>D</sub><sup>25</sup> +22° (c=0.1, MeOH). To a stirred solution of the amine (10 mg, 0.016 mmol) in anisole or thioanisole (80 µL, ≈50 equiv) was added TFA (0.5 mL). The sluggish reaction was monitored by TLC, and after 48 h had proceeded to completion, whereupon 1.0 mL H<sub>2</sub>O was added, and the solution was adjusted to pH10 with 5.0 N NaOH, followed by extraction with CH<sub>2</sub>Cl<sub>2</sub> (3x 5mL). Combined organic layers were dried over Na<sub>2</sub>SO<sub>4</sub>, and evaporated to a residue, which was purified by preparative TLC (5% MeOH:CH<sub>2</sub>Cl<sub>2</sub>) to provide (+)-Staurosporine (49, 6 mg, 70%). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>): δ 9.43 (d, J=7.9 Hz, 1H), 7.94 (8.5, 1H), 7.90 (d, J=7.7 Hz, 1H), 7.49 (app.t, J=7.6 Hz, 1H), 7.43 (app.t, J=7.7 Hz, 1H), 7.37 (app.t, J=7.5 Hz, 1H), 7.33 (app.t, J=7.4 Hz, 1H), 7.30 (d, J=8.0 Hz, 1H), 6.57 (d, J=5.6 Hz, 1H), 6.33 (bs, 1H), 5.05 (d, J=15.8 Hz, 1H), 5.01 (d, J=15.8 Hz, 1H), 3.89 (bs, 1H),

3.42 (s, 3H), 3.37 (d, 3.2H), 2.76 (dd, J=3.9, 14.7 Hz, 1H), 2.41 (bd, J=15.4 Hz, 1H), 2.37 (s, 3H), 1.59 (bs, 1H), 1.57 (s, 3H). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>): δ 173.6, 139.8, 136.7, 132.2, 130.8, 126.6, 125.0, 124.6, 124.2, 123.4, 120.6, 120.0, 119.8, 115.3, 114.1, 106.9, 91.1, 84.2, 80.1, 57.2, 50.4, 45.9, 33.3, 30.3, 30.1. IR (thin film/NaCl): 3316.6 (m), 2925.0 (m), 2850.8 (m), 1678.7 (s), 1636.2 (m), 1584.2 (m), 1457.5 (s), 1352.2 (s), 1316.7 (s), 1281.3 (m), 1115.5 (m), 744.8 (s) cm<sup>-1</sup>. [α]<sub>D</sub><sup>25</sup> +35° (c=0.1, MeOH); natural staurosporine [α]<sub>D</sub><sup>25</sup> +35° (c=1.0, MeOH).

The above description is for the purpose of teaching the person of ordinary skill in the art how to practice the present invention, and it is not intended to detail all those obvious modifications and variations of it which will become apparent to the skilled worker upon reading the description. It is intended, however, that all such obvious modifications and variations be included within the scope of the present invention, which is defined by the following claims. The claims are intended to cover the claimed components and steps in any sequence which is effective to meet the objectives there intended, unless the context specifically indicates the contrary.

**Referenc s**

Fredenhagen, A.; Peter, H.H. *Tetrahedron* 52: 1235 (1996).

Link, J.T., *et al.*, *J. Am. Chem. Soc.* 115: 3782 (1993).

Ootsuka, Y. *et al.* . Jpn. Kokai Tokkyo Koho JP05247054, (1993).

Omura, S. *et al.* *J. Antibiotics* 48: 525 (1995).

McCombie, S.W., *et al.*, *Bioorg. Med. Chem. Lett.* 3: 1537 (1993).

Pirrung, M.C., *et al.*, *J. Org. Chem.* 60: 2112 (1995).


Stolz, B.M., and Wood, J.L., *Tetrahedron Lett.* 36: 8543-8544 (1995).

Stoltz, B.M., and Wood, J.L., *Tetrahedron Lett.* 37: 3929-3930 (1996).

Wood, J.L., *et al.*, *J. Amer. Chem. Soc.* 117: 10413-10414 (1995).

**Claims**


1. A process for the preparation of furanosylated indolocarbazoles of the formula



by reacting an acetal selected from the group consisting of the formulae



and mixtures thereof,  
to produce a glycosylated product of the formula




wherein R is selected from the group consisting of H, C, N, S, O, Cl, Br, I, Si, F, and mixtures thereof.


2. A process according to claim 1 wherein said preparation is carried out under conditions that promote acetal exchange or formation.

-43-


3. A process according to claim 2 wherein said preparation is carried out in the presence of a Bronsted acid or a Lewis acid.
4. A process according to claim 1 wherein camphor sulfonic acid is used as a catalyst and dichloroethane is used as a solvent.
5. A process according to claims 1 to 4 wherein a furanose of the formula



is reacted with DMB-protected K252c to give two products of the formulae




6. A product prepared according to the process of claims 1 to 5.
7. A process for the stereoselective preparation of pyranosylated indolocarbazoles of the formula



-44-


by ring expansion of the corresponding furanosylated indolocarbazole of the formula



wherein R is selected from the group consisting of H, C, N, S, O, Cl, Br, I, Si, F, and mixtures thereof.

8. A process according to claim 7 wherein the ring expansion is carried out in the presence of a Bronsted acid or a Lewis acid.

9. A process according to claims 7 or 8 wherein the ring expansion is carried out in a multistep procedure wherein a compound of the formula




is first reduced with  $\text{LiBH}_4$ , the derived diol is oxidized, and the resulting intermediate compound is then subjected to  $\text{BF}_3 \cdot \text{Et}_2\text{O}$ .

10. A product produced by the process of claims 7, 8, or 9.


11. A compound selected from the group consisting of

-45-




wherein R<sup>1</sup> is selected from the group consisting of H, *t*-Bu, 3,4-dimethoxybenzyl, 4-methoxybenzyl, benzyl, and X and Y are selected from the group consisting of O and H.

12. A compound selected from the group consisting of



wherein  $R^1$  is selected from the group consisting of H, *t*-Bu, 3,4-dimethoxybenzyl, 4-methoxybenzyl, benzyl;  $R^2$  is selected from the group consisting of H and  $CH_3$ ;  $R^3$  is selected from the group consisting of O, NOH, and NOBn;  $R^4$  is selected from the group consisting of H, CHO, and  $CH_3$ ; and X and Y are selected from the group consisting of O and H.


13. A compound selected from the group consisting of



wherein  $R^1$  is selected from the group consisting of, H, *t*-Bu, 3,4-dimethoxybenzyl, 4-methoxybenzyl, benzyl;  $R^2$  is selected from the group consisting of H and  $CH_3$ ;  $R^3$  is selected from the group consisting of O, NOH, and NOBn;  $R^4$  is selected from the group consisting of H, CHO, and  $CH_3$ ; and X and Y are selected from the group consisting of O and H.




14. A compound selected from the group consisting of




-48-

wherein  $R^1$  is selected from the group consisting of H,  $CH_3$ ;  $R^2$  is selected from the group consisting of O, NOH, and NOBn; and  $R^3$  is selected from the group consisting of H, CHO, and  $CH_3$ .


15. A process for preparing tertiary alcohols containing the structural features of compounds selected from the group consisting of



comprising reacting at least one diazo carbonyl compound of the formula



with an allylic alcohol of the formula,



wherein R is selected from the group consisting of H, C, N, S, O, Cl, Br, I, Si, F, and mixtures thereof,


under conditions that produce a carbene or carbenoid intermediate from the diazo compound and for a time sufficient to couple the compounds.

16. A process according to claim 15 wherein the reaction is promoted by thermolysis, photolysis, or a catalyst selected from the group consisting of transition metal catalysts, Bronsted acids, and Lewis acids.


17. A process according to claim 16 wherein the reaction is carried out in the presence of a transition metal catalyst.

-49-


18. A process according to claim 17 wherein the catalyst is  $\text{Rh}_2(\text{OAc})_4$ .
19. A product prepared according to the process of claims 15 to 18.
20. A process for the preparation of an indolocarbazole of the formula



comprising reacting a diazo compound of the formula



with a biindole of the formula




wherein R is selected from the group consisting of H, C, N, S, O, Cl, Br, I, Si, F, and mixtures thereof.


21. A process according to claim 20 wherein the reaction is carried out in the presence of a transition metal catalyst in a solvent capable of solvating the reactants.
22. A process according to claim 21 wherein the coupling reaction is carried out in the presence of a  $\text{Rh}_2(\text{OAc})_4$  catalyst.

-50-

23. A process according to claims 20 to 22 wherein the diazo compound is a diazolaactam and the biindole is a 2,2'-biindole.
24. A product prepared according to the process of claims 20 to 23.
25. A process for the stereoselective preparation of a furanosylated indolocarbazole of the formula



by ring contraction of the corresponding pyranosylated indolocarbazole of the formula



wherein R is selected from the group consisting of H, C, N, S, O, Cl, Br, I, Si, F, and mixtures thereof.

26. A process according to claim 25 wherein the ring contraction is carried out in the presence of a compound capable of oxidizing an alcohol to a ketone and a reagent capable of effecting a benzylic acid-type rearrangement.

-51-

27. A process according to claims 25 or 26 wherein the ring contraction is a single-step procedure employing CuCl as both the oxidant and the rearrangement promoter, and methanol is used as a solvent.

28. A product prepared according to the process of claims 25, 26, or 27.